

Original Article



Effect of Midwifery Students' Continuity of Care on Childbirth Fear and Postpartum Depression: An Individually Randomized Two-Arm Superiority Trial

Elham Jafari¹ , Mojgan Mirghafourvand^{2,3}, Shamsi Abbasalizadeh⁴, Shahla Meedya⁵, Leila Doshmangir⁶, Sakineh Mohammad-Alizadeh-Charandabi^{2,3*}

¹Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran

²Social Determinants of Health Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

³Department of Midwifery, Faculty of Nursing and Midwifery, Tabriz University of Medical Sciences, Tabriz, Iran

⁴Women's Reproductive Health Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

⁵Centre for Nursing and Midwifery Research, School of Nursing and Midwifery, Western Sydney University, Sydney, Australia

⁶Department of Health Policy and Management, Tabriz Health Services Management Research Center, School of Management and Medical Informatics, Tabriz University of Medical Sciences, Tabriz, Iran

ARTICLE INFO

Article History:

Received: September 10, 2025

Revised: December 12, 2025

Accepted: February 16, 2026

ePublished: xx, 2026

Keywords:

Continuity of patient care, Education, Medical, Undergraduate, Iran, Mental health, Midwifery, Pregnancy, Quality of health care, Students

*Corresponding Author:

Sakineh Mohammad-Alizadeh-Charandabi,
Email: mhammadalizadehs@gmail.com

Abstract

Introduction: Midwifery continuity of care may improve maternal mental health. Integrating this model into midwifery education may facilitate broader adoption; however, no trials examined its effects on maternal health. We aimed to assess the effect of a midwifery student's continuity of care program on fear of childbirth and postpartum depression (primary outcomes).

Methods: In this trial, low-risk women at 26–29 weeks of gestation were recruited from Tabriz health centers from November 2022 to February 2023 and assigned to intervention or control groups using stratified block randomization. Twenty-five final-year midwifery students provided continuity of care (each for one to two women) from pregnancy through postpartum for the intervention group; both groups received usual care. Childbirth fear and depression were assessed using the Wijma Delivery Expectancy/Experience Questionnaire and Edinburgh Postnatal Depression Scale at baseline, and 40–50 days postpartum. Analysis of covariance, adjusted for baseline values, parity, and intended hospital type, was used to compare the groups.

Results: All 93 recruited women were analyzed; 50% were primiparous. In the intervention group (n=45), 76% attended ≥6 antenatal sessions, and 64% participated in ≥3 postpartum sessions provided by the students. The intervention group showed lower mean total fear of childbirth (35.8 vs. 75.8, adjusted mean difference (AMD) -43.2; 95% CI: -52.7 to -33.7) and depression scores (5.3 vs. 11.2, AMD -5.9; 95% CI: -8.1 to -3.7) than controls.

Conclusion: The findings support further evaluation and implementation research. However, these findings should be interpreted cautiously considering the single-city design, lack of blinding, and reliance on self-reported outcomes.

Trial registration: Iranian Registry of Clinical Trials, IRCT20100414003706N41, registered prospectively on 30 April 2022, <https://www.irct.ir/search/result?query=IRCT20100414003706N41>.

Introduction

Fear of childbirth (FOC) and postpartum depression (PPD) are major contributors to psychological birth trauma, affecting women's mental health and well-being.¹ A review reported a wide discrepancy in FOC prevalence, ranging from 0.7% to 89% across cultures and countries.² In Iran, reported FOC prevalence ranges from 17% to 89%.³ The global PPD prevalence is 26.3%,⁴ with an estimated 28.7% in Iran.⁵ FOC is associated with increased caesarean section requests, preterm labor, prolonged labor, postpartum depression, and post-traumatic stress disorder.⁶ PPD is also linked to many adverse outcomes, such as poor

psychological health, impaired maternal-infant bonding, relationship difficulties, sexual dysfunction, and reduced quality of life.⁷

Midwifery-led continuity of care (MCOC) is a key framework for enhancing the psychological well-being of women with FOC.⁸⁻¹⁰ A 2024 overview and a 2025 systematic review found that prenatal education, psychoeducation, and counselling may reduce FOC.^{11,12} MCOC may also play a critical role in managing perinatal depression by fostering a trust-based relationship between women and midwives.¹³⁻¹⁵ However, a recent systematic review found limited evidence on the effect of midwifery-led caseload

care on PPD.¹⁶

Midwifery student continuity of care (MSCOC) is a specific form of MCOC in which students provide continuous care to childbearing women. It is part of the midwifery training requirements in some European countries.¹⁷ However, it has been implemented in only one low- and middle-income country, Indonesia.¹⁸ Qualitative and observational studies emphasize that women value emotionally supportive, meaningful relationships with students in MSCOC settings.¹⁹⁻²³

In Iran, midwives' roles in low-risk pregnancy and childbirth are often overlooked²⁴ due to the medicalization of maternity care.²⁵ This medicalization has diminished midwifery authority and hindered a woman-centered approach.²⁶ Despite initiatives to promote physiological birth, including childbirth preparation classes and guidelines,²⁷ the caesarean section remains high at 52%.²⁸ Fear of childbirth, alongside negative experiences, poor communication, and insufficient continuous care, is a key reason for requesting a caesarean.²⁹ Limited evidence from Iran suggests that midwifery-led care improves vaginal birth rates and childbirth satisfaction.^{30,31} Also, a trial in Iran found that a mHealth application providing continuous support reduced FOC.³² To transition to MCOC, reforms in midwifery education, policy, and supportive environments are necessary.¹⁷ Strengthening education and empowering midwifery students in MCOCs are essential for improving service quality and achieving lasting improvements in Iran's midwifery services.^{17,33}

Reviews highlight that 90% of MCOC studies have been conducted in high-income countries,¹⁸ emphasizing the need for research in low- and middle-income countries.³⁴ We also found only limited observational studies and no randomized controlled trials on the effect of MSCOC on maternal health. Therefore, we aimed to implement the MSCOC program during pregnancy, labor, and the postpartum period and to examine its effects on fear of childbirth and postpartum depression.

Methods

Study Design

This study was a two-parallel-arm, superiority, individually randomized controlled trial (RCT). Due to the nature of the intervention, blinding the participants and care providers was not possible. Blinding the outcome assessor was also not possible, as outcomes were self-reported by the participants. This paper adheres fully to the CONSORT guidelines.

Participants

The inclusion criteria for women were singleton pregnancy, gestational age between 26 to 29 weeks, at least six years of education, a history of up to two vaginal births without prior caesarean section, and willingness to have a vaginal birth at participating hospitals. Exclusion criteria included underlying diseases, high-risk pregnancy history, current pregnancy complications, recent stressful events, major fetal abnormalities, lack of smartphone or internet access,

or participation in another trial.

Initially, recruitment was limited to women intending to give birth at two teaching hospitals in Tabriz. Approximately two months after recruitment began, before any data analyses, due to slow enrolment progress, eligibility criteria were expanded to include women intending to deliver at all hospitals with high delivery volumes in the city. One hospital declined permission to participate, resulting in the final addition of one social security hospital and six private/semiprivate hospitals. The expansion was done after approval of the research council, but the registered protocol was not amended to reflect this change. To account for potential differences related to hospital setting, randomization was stratified by hospital type (teaching, social security, and private/semi-private hospitals).

Twenty-five of 28 final-year midwifery students from Tabriz University of Medical Sciences, with written informed consent, participated as MSCOC providers in the intervention group.

The principal investigator (PI, first author, PhD student in midwifery) mentored the students. She is an experienced midwifery instructor and a member of the physiological childbirth training core.

Recruitment

Women from all public health centres in Tabriz, Iran, with at least four potentially eligible pregnant women were recruited. The PI identified potentially eligible women through the national integrated health system (IHS) entitled "SIB". She then contacted the women by phone, provided study details, and invited them to visit the health centre to participate in the trial. After further eligibility screening and providing detailed study information, written informed consent was obtained from eligible women. The women were then recruited into the study and completed the baseline questionnaires.

Sample Size

The sample size was calculated using G*Power software based on an independent samples t-test (two-sided), considering the two pre-specified primary outcomes: postpartum depression (PPD) and fear of childbirth (FOC). A two-sided α of 0.05, 90% power, and a 1:1 allocation ratio were assumed.

Based on a previous Iranian study³⁵, assuming a mean PPD score of 14.9, a standard deviation of 4.0, and a clinically significant reduction of 2.9 points (20% reduction), 40 participants per group were required. After allowing for 15% possible attrition, the final sample size was set at 46 participants per group. For FOC, based on Andaroon et al.³⁶, assuming mean W-DEQ scores of 63.9 and 51.1 in the control and intervention groups, respectively (common SD: 14.1), the required sample size was 33 participants per group. Therefore, the sample size determined based on PPD was considered the final sample size.

Sequence Generation and Randomization

The allocation sequence was generated using a computer

program with randomly varied block sizes of 4 and 6 (1:1 ratio), stratified by parity (nulliparous, multiparous) and intended hospital type for delivery (teaching, social security, private/semiprivate). The sequence was generated by a person not involved in recruitment, randomization, intervention, or data collection. We used a central method to conceal allocation from the investigator (PI) who recruited and assigned participants into the groups. The variable block sizes were concealed from the investigator responsible for participant recruitment and allocation to prevent prediction of upcoming intervention assignments. After participant recruitment and receiving written informed consent, while individuals were completing the baseline questionnaires, the PI sent the participant's name, mobile number, and stratification factors (i.e., parity and intended hospital type for delivery) to the sequence generator via text message, who informed the PI of the woman's group based on the predetermined sequence after completion of baseline questionnaires.

Simple randomization was used to assign women in the intervention group to midwifery students. Each student was responsible for one to two women as primary care providers, and supported one to two others assigned to another student. Supporters were selected based on their relationships with primary care providers, ideally with one being from Tabriz. On the recruitment day, the primary student care provider (and, if possible, her supporter) and the PI attended the centre, where participants in the intervention group were introduced to their student care providers. If the assigned student was inaccessible at the recruitment day, the PI contacted the next student on the randomization list. Pregnant women were assigned to inaccessible students according to a new randomization sequence.

Intervention

Student training and mentoring

Before recruitment, students attended four 4-hour workshops covering communication skills, assertiveness, critical thinking, decision-making, physiological childbirth, evidence-based midwifery, the philosophy of continuity of care, and Iranian national maternal health guidelines. The mentor organized and taught the workshops.

The mentor established a Telegram group for student communication throughout the study to enhance their learning and support. She shared educational content in PDFs with audio and brief videos, and also facilitated weekly online question-and-answer sessions to address students' questions. Students also shared relevant cases related to mothers under their care, such as anonymous test results or audio files. The mentor facilitated discussions and provided comprehensive explanations. Additionally, the mentor privately sent step-by-step training content for each care session to students, two to three days before the scheduled session.

The mentor set dates for subsequent sessions based on the woman's gestational age and coordinated with the student care provider. Students were required to report

session outcomes or cancellations to the mentor.

Continuous care for women in the intervention group

Each woman received continuous care from a midwifery student from recruitment through six weeks postpartum. The care included nine antenatal visits, attendance at birth from the onset of active labor to two hours postpartum, and four postnatal follow-ups. The care began on the recruitment day; during this session, the student introduced herself, outlined the two-way communication process, and conducted the first care session. All sessions included effective communication, addressing women's questions, assessing risk factors, and discussing common complaints. The intervention schedule and core components are summarized in Table 1; detailed session-specific activities, the TIDieR checklist, and the intervention fidelity framework are provided in Supplementary Tables S1, S2, and S3, respectively.

The supporting students participated in one to two video sessions, but not phone sessions. Both students shared contact numbers with the assigned woman for nonemergency questions between 8 a.m. and 11 p.m., with 24-hour access for emergencies. Initially, women contacted the primary provider; if the primary provider was unavailable, they reached out to the supporting student. The mentor's phone number was also provided to the women for emergencies.

The mentor attended all in-person pregnancy sessions, assessed students' clinical and communication skills, and provided feedback. With the women's consent, one to two initial phone/video sessions per student were recorded and shared with the mentor for guidance. The mentor was also on-call for the students. To ensure protocol adherence, the students completed a session report form for the mentor after each session.

Care during labor and birth for the intervention group

To ensure continuity of care during labor and birth, women were asked to notify their primary student care provider when labor symptoms began or upon admission for pregnancy termination. The student would then inform the mentor about the woman's condition. Both remained updated through follow-ups, primarily by phone or occasionally in person, using information provided by the woman or her hospital midwife.

Once active labor began, the student attended to the woman's bedside, keeping her informed of labor progress and providing emotional support. The student also assisted with breathing techniques, position changes, and massage. The student remained with the woman until two hours postpartum. If an emergency caesarean section was required, the student's presence in the operating room had been coordinated, although the staff sometimes restricted it.

Routine care for both groups

All participants had prenatal care at public health centers, with the option to consult private obstetricians or midwives.

Table 1. Key components of the Midwifery Student Continuity of Care (MSCOC) intervention

Phase	Timing	Delivery mode	Key intervention components
Antenatal continuity of care	26–29 weeks until birth	Two face-to-face visits and seven scheduled telephone/video counselling contacts (nine planned antenatal contacts)	Continuous care by the same student midwife; review of medical records; individualized counselling and communication; childbirth preparation and education; assessment of risk factors; scheduled and responsive support based on women's needs.
Labour and birth	Active labour until 2 h postpartum	Face-to-face	Continuous intrapartum support by the woman's primary student care provider when feasible, from active labour until two hours postpartum; emotional support, information on labour progress, and supportive care during labour.
Postpartum continuity of care	12–24 h postpartum until 6 weeks postpartum	One face-to-face visit and three scheduled telephone/video follow-up contacts	Breastfeeding counselling; maternal and newborn care education; postpartum self-care; mental health counselling; reproductive health and family planning counselling; continued individualized support.

National standard care for low-risk pregnancies included eight visits during pregnancy and three visits within the first six weeks postpartum. Also, free childbirth preparation classes, led by midwives not involved in ongoing care, were offered at selected centers²⁷. In teaching hospitals, midwives assist during labor, and female obstetric residents primarily provide care following a medicalized approach³⁷. In other hospitals, midwives work under an obstetrician's supervision with limited autonomy. Private hospitals offer more respectful care, but continuous midwife support during labor and pharmacological pain relief incur additional costs²⁸.

Outcomes

Among the prespecified primary outcomes, FOC and PPD assessed at 40–50 days postpartum are reported in the current manuscript. Due to space constraints, results of childbirth experience, the third prespecified primary outcome, are reported separately in a companion publication, in which the complete trial protocol is provided as supplementary material.³⁸ Secondary outcomes included FOC and PPD assessed at 35–36 weeks of pregnancy and FOC subscale scores at both assessment time points. No hierarchy was prespecified among the primary outcomes, and no multiplicity adjustment was applied; each primary outcome was interpreted separately.

Data Collection Tools

The Wijma Delivery Expectancy/Experience Questionnaire (W-DEQ) was used to assess FOC. It has two versions, each comprising 33 six-point items (scored 0–5) and six subscales. W-DEQ-A assesses women's prenatal expectancies regarding childbirth, while W-DEQ-B evaluates their recent childbirth experiences.³⁹ Previous studies have established the validity and reliability of the W-DEQ in the Iranian context.⁴⁰ In the present study, Cronbach's alpha values were calculated to assess internal consistency and were 0.97 for W-DEQ-A and 0.96 for W-DEQ-B. Alpha values for the subscales ranged from 0.70 to 0.98 for version A and from 0.67 to 0.96 for version B.

The Edinburgh Postnatal Depression Scale (EPDS), a valid scale to assess depression symptoms during pregnancy and postpartum, was used to assess depression. It consists of 10 four-point Likert items (scored 0–3).⁴¹ Its validity and reliability have been confirmed in Iran, with a Cronbach's alpha of 0.77 and an ICC of 0.80 at 6–8 weeks postpartum.⁴² In our study, Cronbach's alpha was 0.89 at 35–36 weeks of

pregnancy and 0.92 at 40–50 days postpartum.

A demographic and obstetric data form, developed for this study, was used to collect baseline data. Experts confirmed its content validity.

Data Analysis

Data were analyzed using IBM SPSS Statistics for Windows, Version 24.0 (IBM Corp., Armonk, NY, USA). A range check and an independent review of 10% of randomly selected cases were conducted to verify the accuracy of data entry. Questionnaires were reviewed immediately after completion, and participants were asked to complete any missing responses.

Data analysis was performed after completion of data collection in accordance with the intention-to-treat principle. Separate ANCOVA models were used to compare post-intervention outcomes between the MSCOC and control groups. Baseline outcome values were included as covariates, while intervention group, parity, and hospital type were entered as fixed factors, consistent with the stratified randomization approach. We used complete-case analyses with no imputation for the missing values for the four participants (two per group) at the 35–36-week assessment (secondary outcomes).

The assumptions of ANCOVA were assessed, including normal distribution of dependent variables and linearity relation between baseline and post-intervention scores at each intervention group, homogeneity of regression slopes, normality of residuals, and homogeneity of variance. All assumptions were considered acceptable for the analyses performed. A two-sided $P < 0.05$ was considered statistically significant.

Exploratory post-randomization subgroup analyses were conducted based on student attendance during labor and delivery within the MSCOC group; as well as comparison of the MSCOC group without student attendance during labor and delivery with the control group.

Results

Recruitment occurred from November 2022 to February 2023, with follow-up completed in July 2023. Of the 60 health centres in Tabriz, 27 met the sampling criteria, and 1–6 women were recruited from each centre. Of the 280 women assessed for eligibility, 93 eligible women were randomly allocated to either the MSCOC group ($n = 45$) or the control group ($n = 48$). All randomized women completed postpartum follow-up (primary outcome

assessment). However, four participants (two from each group) were not assessed at 35–36 weeks of pregnancy. All randomized participants were analyzed according to the intention-to-treat principle (Figure 1).

Baseline demographic and obstetric characteristics of the participants, including stratification variables (parity and intended hospital type for delivery) by study groups, are presented descriptively in Table 2. The mean age of participants was 26.9 years (SD 4.6); 54% had a diploma or university education, and 50% were primiparous.

Twenty-eight women (62%) in the MSCOC group and 35 women (73%) in the control group received prenatal care from a private obstetrician, in addition to routine care at the public health centers. In the MSCOC group, 3 women (7%) attended childbirth preparation classes, compared to 7 women (15%) in the control group. During labor and birth, 2 women (7%) in the MSCOC group were accompanied by private midwives and 10 (37%) by family members. In the control group, the corresponding figures were 4 (10%) and 8 (21%). Women attended a mean of 7.3 antenatal sessions (SD = 2.8; range: 1–9) and 2.8 postnatal sessions (SD = 1.3; range: 0–4). Details of the protocol adherence in the intervention group are presented in Table 3.

At baseline, the mean total FOC score was numerically higher in the MSCOC group than in the control group (62.1 vs. 54.5). Post-intervention comparisons were performed using ANCOVA models adjusted for baseline FOC score, parity, and hospital type. The MSCOC group had significantly lower mean total FOC scores at 35–36 weeks of pregnancy (31.0 vs. 75.3; adjusted mean difference (AMD)

Table 2. Baseline characteristics of participants by study groups

Characteristics	MSCOC (n=45)	Control (n=48)
Age (years)	27.1 [6.3]	26.7 [6.6]
Education (Diploma or college)	22 (48.9)	28 (58.3)
Job (Unemployed)	44 (97.8)	41 (85.4)
Adequacy of family income		
Not enough at all	3 (6.7)	3 (6.3)
Relatively enough	38 (84.4)	38 (79.1)
Quite enough	4 (8.9)	7 (14.6)
History of infertility (Yes)	6 (13.3)	2 (4.2)
Planned pregnancy	33 (73.4)	33 (68.8)
Primiparous	22 (46.8)	25 (53.2)
Pregnancy interval from previous birth (years)*	7.7 [3.4]	6.2 [3.2]
Gestational age at recruitment (weeks)	27.5 [1.3]	27.6 [1.2]
Sex of the fetus (Female)	21 (46.7)	23 (47.9)
Satisfaction from previous childbirth*		
Very satisfied or satisfied	11 (47.9)	10 (43.5)
Neither satisfied nor dissatisfied	5 (21.7)	8 (34.8)
Dissatisfied or very dissatisfied	7 (30.4)	5 (21.7)
Intended hospital type for delivery		
Teaching	18 (40)	20 (41.7)
Social security	9 (20)	7 (14.6)
Private/semiprivate	18 (40)	21 (43.7)

MSCOC: Midwifery student continuity of care

The data are presented as number (%) or mean [SD]

* Among individuals who had a previous birth, n=23 in each group




Figure 1. CONSORT Flowchart of Participant Allocation

-47.8, 95% CI -54.6 to -40.9) and at 40–50 days postpartum (35.8 vs. 75.8; AMD -43.2, 95% CI -52.7 to -33.7) compared with the control group. (Table 4). At both post-intervention assessments, all FOC subscale scores were lower in the MSCOC group than in the control group (Table S4).

The mean depression score at baseline was 11.2 in both the intervention and control groups. However, significantly lower scores were observed in the intervention group at both postintervention follow-ups: gestational weeks 35–36 (7.1 vs. 13.5; AMD -6.4; 95% CI -8.2 to -4.5) and 40–50 days postpartum (5.3 vs. 11.2; AMD: -5.9; 95% CI: -8.1 to -3.7).

There were no statistically significant differences between subgroups of the MSCOC group with and without student attendance during labor and delivery in terms of frequency of primiparity, main antenatal care provider, or type of delivery. However, the subgroup with student attendance had significantly higher frequency of attendance at six or more antenatal sessions (95% vs 58%, $P=0.005$) and at three or more postpartum sessions (95% vs 38%, $P<0.001$). They also had a higher frequency of delivering at a teaching hospital (52% vs 29%, $P=0.03$) (Table S5).

As an exploratory analysis, we examined outcomes according to student attendance during labor and delivery. At 35–36 weeks of pregnancy, no statistically significant differences were observed between the two subgroups of the MSCOC group, with and without student attendance during labor and delivery, in the mean total FOC, FOC subscales, or the depression scores. However, at 40–50 days postpartum, participants with student attendance had lower loneliness and fear subscale scores. Total FOC and depression scores were also numerically lower (by 12 and 2.5 points, respectively), but the differences were not statistically significant (Table S6).

At 40–50 days postpartum, no statistically significant

differences were observed between the subgroup without student attendance during labor and delivery and the control group in the mean scores for total FOC and PPD scores. The group without student attendance had significantly lower mean scores than the control group on two FOC subscales: the lack of self-efficacy and the concerns for the child (S3).

Discussion

In this study, we introduced the MSCOC model in Iran, a novel approach in a context where MCOC has not been widely implemented. The MSCOC group showed lower post-intervention FOC severity and PPD symptoms than the control group. The observed reduction in W-DEQ scores exceeded commonly used thresholds for clinically relevant fear of childbirth, suggesting that the intervention effect may have clinical relevance in addition to statistical significance. Moreover, the reduction in EPDS scores moved participants toward a lower-risk range based on established screening thresholds for postpartum depressive symptoms.

The findings are consistent with previous Swedish research highlighting the benefits of MCOC for women experiencing FOC.^{43–45} Another qualitative study reported that women experienced reduced stress and increased autonomy through MCOC.⁴⁵ Additionally, an experimental study indicated that MCOC offers valuable opportunities for decision-making participation, and a sense of control for women with FOC.⁴⁴ The establishment of a mutually trusting relationship between women and midwives can reduce their fear.⁴⁶ A prior Iranian study found that the childbirth preparation courses approved by the Ministry of Health did not significantly reduce FOC.⁴⁷ While these courses enhance knowledge, their effectiveness in

Table 3. Protocol adherence in the intervention group

Category	Reported reason for non-complete adherence
34 women (76%) attended ≥ 6 antenatal sessions	- Lack of interest (3 cases) - Inadequate follow-up by the student (3 cases)
29 women (64%) attended ≥ 3 postpartum sessions	- Spouse's disapproval (2 cases) - Sudden death of close relatives, opting out of vaginal birth based on obstetrician advice, and preterm birth (each, one case)
In 21 cases (47%), the student attended during Labor and delivery (In 20 out of 21 cases, the primary student care provider was present)	- Women not informing students (20 cases) due to emergency childbirth, lack of phone access in the delivery room (5 cases), embarrassment to call due to midnight childbirth (2 cases). The 13 other cases that did not provide specific reasons were mostly those who had attended five or fewer antenatal sessions (10 cases). - Inadequate follow-up by the student (4 cases).

Table 4. Comparison of childbirth fear and depression scores between study groups

Outcomes	Baseline		35–36 weeks of pregnancy		40–50 days postpartum	
	MSCOC n=45	Control n=48	MSCOC n=43	Control n=46	MSCOC n=45	Control n=48
Total Childbirth fear (0–165)	62.1 (19.9)	54.5 (27.8)	31.0 (18.6)	75.3 (21.1)	35.8 (23.7)	75.8 (26.1)
AMD (95% CI)*	7.0 (-3.0 to 16.9)		-47.8 (-54.6 to -40.9)**		-43.2 (-52.7 to -33.7)**	
Depression (0–30)	11.2 (6.2)	11.2 (6.0)	7.1 (4.2)	13.5 (5.8)	5.3 (4.6)	11.2 (6.6)
AMD (95% CI)*	-0.2 (-2.7 to 2.3)		-6.4 (-8.2 to -4.5)**		-5.9 (-8.1 to -3.7)**	

The data are presented as mean (standard deviation) unless otherwise specified.

MSCOC: Midwifery Student Continuity of Care; AMD: adjusted mean difference between the groups; CI: confidence interval

The Wijma Delivery Expectancy/Experience Questionnaire (W-DEQ) was used to assess fear of childbirth, and the Edinburgh Postnatal Depression Scale (EPDS) was utilized to assess depression symptoms. Higher scores indicate greater levels of fear or depression.

* ANCOVA was used, adjusted for parity and hospital type for the baseline comparisons, and adjusted for parity, hospital type, and baseline values for post-intervention comparisons ** $P<0.001$

promoting a positive childbirth experience is limited.⁴⁸ This is likely due to the lack of individualized, continuity-based care and insufficient focus on FOC.

Although we found no studies examining the effect of MSCOC on FOC, existing literature suggests that MSCOC shares several characteristics with MCOC that may help alleviate FOC. This model has been shown to enhance women's empowerment through supportive care, strengthening the partnership between women and students, and providing evidence-based care.¹⁹⁻²¹ Qualitative studies indicated that women perceive the student midwife's attendance in labor and delivery as essential.^{22,23} They view the student midwife as their advocate, helping them express their needs, expectations, and concerns, fostering calm and security, and enabling greater focus on the childbirth process.¹⁹ Our exploratory subgroup analysis showed that participants with student attendance during labor had lower FOC subscale scores related to fear and loneliness; however, these findings represent associations and should not be interpreted as causal effects of student attendance. Students were more likely to attend labor for women who had participated in most prenatal and postpartum sessions, consistent with findings from a Swedish study.⁴⁴ This pattern may reflect stronger engagement and relationship-building through repeated interactions, although differences in motivation or adherence may also have contributed. Since student attendance during labor was not randomized, these findings should be interpreted as exploratory associations rather than causal effects. Notably, despite the generally negative childbirth experiences reported in teaching hospitals in our context⁴⁹⁻⁵¹, a higher proportion of women in the MSCOC group gave birth in teaching hospitals. The availability of a familiar student midwife and continuous support may have contributed to women's willingness to give birth in these settings; however, differences in hospital type between subgroups may also have influenced the observed findings. Therefore, these subgroup results should be considered hypothesis-generating and interpreted cautiously.

While previous studies did not directly address the effect of MSCOC on PPD, our findings align with research suggesting that psychosocial interventions based on continuous care can significantly reduce PPD risk.^{52,53} Our results are consistent with a Thai study, which found that midwife and family support reduced PPD symptoms.⁵⁴ Studies indicate that brief interventions, such as digital content or limited midwife contacts, do not significantly affect PPD,^{55,56} but MCOC enhances midwives' role in perinatal mental health by providing sufficient time and frequent interactions in a supportive environment.⁸ However, midwives face challenges due to insufficient training in perinatal mental health.⁵⁷ Specialized training in relevant psychosocial skills improves midwives' competence and confidence, positively impacting perinatal mental health, especially in low- and middle-income countries.⁸ In this study, MSCOC providers received such training from a mentor with expertise in perinatal mental health and midwifery-led counseling, which likely contributed

to the observed positive effects.

The study also highlights the challenges related to student attendance during labor and delivery, with only 50% attendance compared to 93% in Australia.²⁰ This discrepancy can be attributed to the voluntary nature of student participation in this study, limited logistical support, and the lack of mandatory educational requirements. The Australian Nursing and Midwifery Council now mandates student involvement during labor as part of the MSCOC requirements.²⁰ However, we observed that students were more likely to attend labor for women who had participated in most prenatal sessions, which aligns with a Swedish study.⁴⁴ This reinforces the importance of building strong, trusting relationships between midwives and women.^{15,58}

In this trial, strict adherence to random sequence generation and central allocation concealment minimized selection bias. Although blinding of participants, student care providers, and outcome assessors was not feasible, efforts were made to reduce potential performance and detection bias by withholding study outcomes from participants and care providers and emphasizing accurate questionnaire completion. The minimal loss to follow-up reduced the risk of attrition bias. Active engagement between the mentor and students likely bolstered adherence to the intervention. Moreover, enrolling participants from all eligible urban health centers in the capital city may have enhanced the generalizability of our findings. Given that most maternity indices in this city (the center of Iran's fifth-largest province) align with the mean urban indices, our findings may be relevant to similar urban settings across Iran.

The study also highlighted the feasibility of integrating MSCOC within our midwifery education system; However, several limitations should be considered.

The intervention was supervised by a single qualified mentor who guided 25 students. As this project was conducted as part of her doctoral thesis, her high motivation and commitment may not be fully representative of larger-scale implementation involving multiple mentors. Due to the mentor's workload and participants' limited internet access, a virtual group could not be established, and some intervention sessions were conducted by telephone. We did not systematically record the frequency of phone calls or text messages exchanged between women and student midwives; therefore, the contribution of these unscheduled supportive interactions to the overall intervention effect remains unclear.

Due to the nature of the intervention, blinding of participants, student care providers, and the mentor who also collected the data was not feasible. Because both primary outcomes were self-reported by participants, the possibility of performance and detection bias cannot be completely excluded.

Although the intervention was delivered by multiple students, clustering by student was not accounted for in the analyses because each student provided care to only one or two women as the primary care provider. Therefore, some residual provider-level variability cannot be completely

excluded. We conducted no multiplicity adjustment due to the three pre-specified primary outcomes, which could lead to increasing type I error. However, because in all primary outcome comparisons P-values were less than 0.001, with Bonferroni correction, the errors will be less than 0.003.

Potential co-interventions may also have influenced maternal outcomes. Although private obstetrician care, childbirth preparation classes, and labor companionship were documented, residual confounding from these factors cannot be completely excluded. Future studies with larger sample sizes should further evaluate their potential influence.

The broad exclusion criteria, smartphone/internet requirements for intervention delivery, and recruitment from urban health centers may limit the generalizability of findings to rural populations, women with limited digital access, and higher-risk pregnancies. Additionally, the limited representation of women from higher socioeconomic backgrounds may restrict applicability to more diverse populations.

The lack of financial incentives or educational requirements for student participation, along with insufficient support for project implementation at hospitals, may have affected student motivation and adherence.

Baseline differences in FOC scores between groups may have influenced the observed changes. Although baseline-adjusted ANCOVA models were used to account for initial differences, residual confounding and regression to the mean cannot be completely excluded.

As PPD and FOC can have negative long-term effects on maternal physical and psychological health, infants' emotional and cognitive development,⁷ and family well-being,⁵⁹ the results of this study offer promising potential for improving family health outcomes. The program offers dual benefits by providing mothers with personalized, woman-centered care while enabling student midwives to enhance their midwifery and interpersonal skills, boost confidence, and improve competence through practical experience.¹⁷ However, there are ambiguities regarding the pedagogical intent and the definition and measurement of learning outcomes for the MSCOC.¹⁷ Given these ambiguities and our study's limitations, we recommend further research across various academic centers. Such research should include action research to explore MSCOC's benefits, challenges, and implementation strategies. It seems that incorporating MSCOC into student curricula, mandating mentoring for academic midwives, involving clinical midwives for student support, and providing facilities could enhance MSCOC implementation and outcomes.

Conclusion

Our findings highlight the potential for integrating MSCOC into the midwifery education system in Iran. The study suggests that MSCOC reduces fear of childbirth and postpartum depression symptoms in women. These findings could motivate further research to explore the advantages, challenges, and potential national adoption of the program, particularly in low- and middle-income

countries.

Acknowledgments

The authors express their gratitude to the Vice-Chancellor for Research of Tabriz

University of Medical Sciences for financial support, and sincere appreciation to the midwifery students who participated in this trial as care providers and to the women who participated in the study. We also acknowledge the valuable support provided by the Clinical Research Development Unit of Taleghani Hospital, Tabriz University of Medical Sciences, Tabriz, Iran.

Authors' Contribution

Conceptualization: Elham Jafari, Mojgan Mirghafourvand, Shamsi Abbasalizadeh, Shahla Meedya, Leila Doshmangir, Sakineh Mohammad-Alizadeh-Charandabi

Data curation: Elham Jafari, Sakineh Mohammad-Alizadeh-Charandabi

Formal analysis: Elham Jafari, Sakineh Mohammad-Alizadeh-Charandabi

Funding acquisition: Sakineh Mohammad-Alizadeh-Charandabi

Investigation: Elham Jafari

Methodology: Elham Jafari, Mojgan Mirghafourvand, Shamsi Abbasalizadeh, Shahla Meedya, Leila Doshmangir, Sakineh Mohammad-Alizadeh-Charandabi

Project Administration: Elham Jafari, Sakineh Mohammad-Alizadeh-Charandabi

Resources: Sakineh Mohammad-Alizadeh-Charandabi

Software: Elham Jafari

Supervision: Sakineh Mohammad-Alizadeh-Charandabi

Validation: Sakineh Mohammad-Alizadeh-Charandabi

Visualization: Elham Jafari, Sakineh Mohammad-Alizadeh-Charandabi

Writing—original draft: Elham Jafari

Writing—review & editing: Elham Jafari, Mojgan Mirghafourvand, Shamsi Abbasalizadeh, Shahla Meedya, Leila Doshmangir, Sakineh Mohammad-Alizadeh-Charandabi

Competing Interests

The authors declare that they have no competing interests.

Consent for Publication

Not applicable.

Data Availability Statement

The corresponding author will make deidentified participant datasets available for research purposes to researchers affiliated with academic institutions and others upon reasonable request immediately after the publication of the results.

Ethics Approval

This study received scientific approval (Grant no: 69665) and ethical approval (Ethical code: IR.TBZMED.REC.1401.094) from Tabriz University of Medical Sciences, Tabriz, Iran.

All participants provided written informed consent before enrolling in the study. The study was conducted in accordance with relevant guidelines and regulations.

Funding

The study received funding from the Vice-Chancellor for Research of the Tabriz University of Medical Sciences. The funding center did not have any involvement in the study design, participant recruitment, data collection, analysis, interpretation, manuscript writing, or the decision of where to submit the manuscript.

Abbreviations

AMD: Adjusted mean difference, EPDS: Edinburgh Postnatal Depression Scale, FOC: fear of childbirth, MCOC: Midwifery continuity of care, MSCOC: Midwifery Student Continuity of Care,

PPD: Postpartum depression, RCT: Randomized controlled trial, W-DEQ: Wijma Delivery Expectancy/Experience Questionnaire.

Supplementary files

Supplementary file 1 contains Table S1-S6.

References

- Horsch A, Garthus-Niegel S, Ayers S, Chandra P, Hartmann K, Vaisbuch E, et al. Childbirth-related posttraumatic stress disorder: definition, risk factors, pathophysiology, diagnosis, prevention, and treatment. *Am J Obstet Gynecol* 2024;230(35): S1116-27. doi:10.1016/j.ajog.2023.09.089
- Dai J, Shi Y, Guo L, Wang Y, Kong X. Discrepancy in parental fear of childbirth: a scoping review. *Midwifery* 2023;126:103830. doi:10.1016/j.midw.2023.103830
- Moradi M, Niazi A, Mazloumi E. Prevalence and causes related to fear of vaginal delivery in Iran: a systematic review. *Iran J Nurs Res* 2022;17(1):43-53. doi:10.22034/ijnr.17.1.43
- Al-Abri K, Edge D, Armitage CJ. Prevalence and correlates of perinatal depression. *Soc Psychiatry Psychiatr Epidemiol* 2023;58(11):1581-90. doi:10.1007/s00127-022-02386-9
- Veisani Y, Sayehmiri K. Prevalence of postpartum depression in Iran-a systematic review and meta-analysis. *Iran J Obstet Gynecol Infertil* 2012;15(14):21-9. doi:10.22038/ijogi.2012.5689
- Imakawa CS, Nadai MN, Reis M, Quintana SM, Moises ECD. Is it necessary to evaluate fear of childbirth in pregnant women? A scoping review. *Rev Bras Ginecol Obstet* 2022;44(7):692-700. doi:10.1055/s-0042-1751062
- Slomian J, Honvo G, Emonts P, Reginster JY, Bruyère O. Consequences of maternal postpartum depression: a systematic review of maternal and infant outcomes. *Womens Health (Lond)* 2019;15:1745506519844044. doi:10.1177/1745506519844044
- Coates D, Foureur M. The role and competence of midwives in supporting women with mental health concerns during the perinatal period: a scoping review. *Health Soc Care Community* 2019;27(4): e389-405. doi:10.1111/hsc.12740
- Fenwick J, Toohill J, Slavin V, Creedy DK, Gamble J. Improving psychoeducation for women fearful of childbirth: evaluation of a research translation project. *Women Birth* 2018;31(1):1-9. doi:10.1016/j.wombi.2017.06.004
- Hildingsson I, Rubertsson C, Karlström A, Haines H. Caseload midwifery for women with fear of birth is a feasible option. *Sex Reprod Healthc* 2018;16:50-5. doi:10.1016/j.srhc.2018.02.006
- Alizadeh-Dibazari Z, Maghalain M, Mirghafourvand M. The effect of non-pharmacological prenatal interventions on fear of childbirth: an overview of systematic reviews and meta-analysis. *BMC Psychiatry* 2024;24(1):415. doi:10.1186/s12888-024-05870-5
- Zhou J, Zhu Z, Li R, Guo X, Li D. Comparative efficacy of non-pharmacological interventions on fear of childbirth for pregnant women: a systematic review and network meta-analysis. *Front Psychol* 2025;16:1530311. doi:10.3389/fpsyg.2025.1530311
- Morlans-Lanau M, González-Vives ML, Rodríguez-Quiroga A, Casbas MM, Klugarová J, Klugar M. Establishing midwife-led continuity of care interventions in perinatal mental health in high-risk pregnancies: a best practice implementation project. *JBIM Evid Implement* 2022;20(S1): S49-58. doi:10.1097/xe.0000000000000324
- Ravaldi C, Mosconi L, Crescioli G, Lombardo G, Russo I, Morese A, et al. Are midwives trained to recognise perinatal depression symptoms? Results of MAMA (MAternal Mood Assessment) cross-sectional survey in Italy. *Arch Womens Ment Health* 2024;27(4):567-76. doi:10.1007/s00737-024-01439-z
- Cibralic S, Pickup W, Diaz AM, Kohlhoff J, Karlov L, Stylianakis A, et al. The impact of midwifery continuity of care on maternal mental health: a narrative systematic review. *Midwifery* 2023;116:103546. doi:10.1016/j.midw.2022.103546
- Wassén L, Borgström Bolmsjö B, Frantz S, Hagman A, Lindroth M, Rubertsson C, et al. Child and maternal benefits and risks of caseload midwifery - a systematic review and meta-analysis. *BMC Pregnancy Childbirth* 2023;23(1):663. doi:10.1186/s12884-023-05967-x
- Hainsworth N, Dowse E, Ebert L, Foureur M. 'Continuity of Care Experiences' within pre-registration midwifery education programs: a scoping review. *Women Birth* 2021;34(6):514-30. doi:10.1016/j.wombi.2020.12.003
- Bradford BF, Wilson AN, Portela A, McConville F, Fernandez Turienzo C, Homer CS. Midwifery continuity of care: a scoping review of where, how, by whom and for whom? *PLOS Glob Public Health* 2022;2(10): e0000935. doi:10.1371/journal.pgph.0000935
- Aune I, Haugen K, Holst-Jenson M, Lund-Tyvoll I. Women's experiences of continuity of care from student midwives - a qualitative study from Norway. *Sex Reprod Healthc* 2023;35:100814. doi:10.1016/j.srhc.2023.100814
- Tickle N, Gamble J, Creedy DK. Women's reports of satisfaction and respect with continuity of care experiences by students: findings from a routine, online survey. *Women Birth* 2021;34(6): e592-8. doi:10.1016/j.wombi.2020.11.004
- Stulz V, Elmir DR, Reilly H. Evaluation of a student-led midwifery group practice: a woman's perspective. *Midwifery* 2020;86:102691. doi:10.1016/j.midw.2020.102691
- Tickle N, Sidebotham M, Fenwick J, Gamble J. Women's experiences of having a bachelor of midwifery student provide continuity of care. *Women Birth* 2016;29(3):245-51. doi:10.1016/j.wombi.2015.11.002
- Jefford E, Nolan SJ, Sansone H, Provost SC. 'A match made in midwifery': women's perceptions of student midwife partnerships. *Women Birth* 2020;33(2):193-8. doi:10.1016/j.wombi.2018.11.018
- Hajiesmaello M, Hajian S, Riazi H, Alavi Majd H, Yavarian R. Challenges facing clinical midwifery education in Iran. *BMC Med Educ* 2022;22(1):407. doi:10.1186/s12909-022-03485-6
- Sandall J, Soltani H, Gates S, Shennan A, Devane D. Midwife-led continuity models versus other models of care for childbearing women. *Cochrane Database Syst Rev* 2016;4(4): CD004667. doi:10.1002/14651858.CD004667.pub5
- Sedigh Mobarakabadi S, Mirzaei Najmabadi K, Ghazi Tabatabaie M. Ambivalence towards childbirth in a medicalized context: a qualitative inquiry among Iranian mothers. *Iran Red Crescent Med J* 2015;17(3): e24262. doi:10.5812/ircmj.24262
- Alizadeh-Dibazari Z, Abbasalizadeh F, Mohammad-Alizadeh-Charandabi S, Jahanfar S, Mirghafourvand M. Childbirth preparation and its facilitating and inhibiting factors from the perspectives of pregnant and postpartum women in Tabriz-Iran: a qualitative study. *Reprod Health* 2024;21(1):106. doi:10.1186/s12978-024-01844-8
- Pourshirazi M, Heidarzadeh M, Taheri M, Esmaily H, Babaey F, Talkhi N, et al. Cesarean delivery in Iran: a population-based analysis using the Robson classification system. *BMC Pregnancy Childbirth* 2022;22(1):185. doi:10.1186/s12884-022-04517-1
- Shirzad M, Shakibazadeh E, Hajimiri K, Betran AP, Jahanfar S, Bohren MA, et al. Prevalence of and reasons for women's, family members', and health professionals' preferences for cesarean section in Iran: a mixed-methods systematic review. *Reprod Health* 2021;18(1):3. doi:10.1186/s12978-020-01047-x
- Bagheri A, Simbar M, Samimi M, Nahidi F, Alavimajd H, Sadat Z. Comparing the implications of midwifery-led care and standard model on maternal and neonatal outcomes during pregnancy, childbirth and postpartum. *J Midwifery Reprod Health* 2021;9(3):2863-72. doi:10.22038/jmrh.2021.55639.1678
- Shahinfar S, Abedi P, Najafian M, Abbaspoor Z, Mohammadi E, Alianmoghadam N, et al. Effect of continuity of team midwifery care on maternal and neonatal outcomes: a quasi-experimental study in Iran. *Sci Rep* 2024;14(1):22819. doi:10.1038/s41598-024-73751-8

32. Khademioore S, Ebrahimi E, Khosravi A, Movahedi S. The effect of an mHealth application based on continuous support and education on fear of childbirth, self-efficacy, and birth mode in primiparous women: a randomized controlled trial. *PLoS One* 2023;18(11): e0293815. doi:10.1371/journal.pone.0293815
33. Khosravi S, Babaey F, Abedi P, Mazaheri Kalahroodi Z, Hajimirzaie SS. Strategies to improve the quality of midwifery care and developing midwife-centered care in Iran: analyzing the attitudes of midwifery experts. *BMC Pregnancy Childbirth* 2022;22(1):40. doi:10.1186/s12884-022-04379-7
34. Sandall J, Fernandez Turienzo C, Devane D, Soltani H, Gillespie P, Gates S, et al. Midwife continuity of care models versus other models of care for childbearing women. *Cochrane Database Syst Rev* 2024;4(4): CD004667. doi:10.1002/14651858.CD004667.pub6
35. Afshari P, Tadayon M, Abedi P, Yazdizadeh S. Prevalence and related factors of postpartum depression among reproductive aged women in Ahvaz, Iran. *Health Care Women Int* 2020;41(3):255-65. doi:10.1080/07399332.2019.1578779
36. Andaroon N, Kordi M, Kimiaei SA, Esmaeily H. The effect of individual counseling program by a midwife on fear of childbirth in primiparous women. *J Educ Health Promot* 2017;6:97. doi:10.4103/jehp.jehp_172_16
37. Abdolalipour S, Abbasalizadeh S, Mohammad-Alizadeh-Charandabi S, Abbasalizadeh F, Jahanfar S, Raphi F, et al. Effect of implementation of the WHO intrapartum care model on maternal and neonatal outcomes: a randomized control trial. *BMC Pregnancy Childbirth* 2024;24(1):283. doi:10.1186/s12884-024-06449-4
38. Jafari E, Mirghafourvand M, Abbasalizadeh S, Meedya S, Doshmangir L, Mohammad-Alizadeh-Charandabi S. Effect of midwifery students' continuity of care program on women's experiences of maternity care: a randomised controlled trial. *PLoS One* 2026;21(7): e0353118. doi:10.1371/journal.pone.0353118
39. Wijma K, Wijma B, Zar M. Psychometric aspects of the W-DEQ; a new questionnaire for the measurement of fear of childbirth. *J Psychosom Obstet Gynaecol* 1998;19(2):84-97. doi:10.3109/01674829809048501
40. Mortazavi F. Validity and reliability of the Farsi version of Wijma delivery expectancy questionnaire: an exploratory and confirmatory factor analysis. *Electron Physician* 2017;9(6):4606-15. doi:10.19082/4606
41. Levis B, Negeri Z, Sun Y, Benedetti A, Thombs BD. Accuracy of the Edinburgh Postnatal Depression Scale (EPDS) for screening to detect major depression among pregnant and postpartum women: systematic review and meta-analysis of individual participant data. *BMJ* 2020;371: m4022. doi:10.1136/bmj.m4022
42. Montazeri A, Torkan B, Omidvari S. The Edinburgh Postnatal Depression Scale (EPDS): translation and validation study of the Iranian version. *BMC Psychiatry* 2007;7:11. doi:10.1186/1471-244x-7-11
43. Fahlbeck H, Hildingsson I, Larsson B, Johansson M. Women's interest in midwifery continuity of care during and after pregnancy and childbirth in Sweden: "as a matter of course". *J Midwifery Womens Health* 2026;71(1):95-103. doi:10.1111/jmwh.70035
44. Hildingsson I, Rubertsson C, Karlström A, Haines H. A known midwife can make a difference for women with fear of childbirth- birth outcome and women's experiences of intrapartum care. *Sex Reprod Healthc* 2019;21:33-8. doi:10.1016/j.srhc.2019.06.004
45. Nääs KL, Johansson M, Wiklund I, Hildingsson I. Women's experiences of participating in a digital continuity of care model designed for fear of birth in a rural setting. *Sex Reprod Healthc* 2025;44:101081. doi:10.1016/j.srhc.2025.101081
46. Larsson B, Rubertsson C, Hildingsson I. A modified caseload midwifery model for women with fear of birth, women's and midwives' experiences: a qualitative study. *Sex Reprod Healthc* 2020;24:100504. doi:10.1016/j.srhc.2020.100504
47. Mousavi SR, Amiri-Farahani L, Haghani S, Pezaro S. Comparing the effect of childbirth preparation courses delivered both in-person and via social media on pregnancy experience, fear of childbirth, birth preference and mode of birth in pregnant Iranian women: a quasi-experimental study. *PLoS One* 2022;17(8): e0272613. doi:10.1371/journal.pone.0272613
48. Mohaghegh Z, Javadnoori M, Najafian M, Montazeri S, Abedi P, Kazemnejad Leyli E, et al. Implementation of birth plans integrated into childbirth preparation classes for vaginal birth: a qualitative study of women, their husbands and clinicians' perspectives in Iran. *BMC Pregnancy Childbirth* 2022;22(1):969. doi:10.1186/s12884-022-05305-7
49. Ghanbari-Homayi S, Fardiazar Z, Meedya S, Mohammad-Alizadeh-Charandabi S, Asghari-Jafarabadi M, Mohammadi E, et al. Predictors of traumatic birth experience among a group of Iranian primipara women: a cross sectional study. *BMC Pregnancy Childbirth* 2019;19(1):182. doi:10.1186/s12884-019-2333-4
50. Hajizadeh K, Vaezi M, Meedya S, Mohammad-Alizadeh-Charandabi S, Mirghafourvand M. Prevalence and predictors of perceived disrespectful maternity care in postpartum Iranian women: a cross-sectional study. *BMC Pregnancy Childbirth* 2020;20(1):463. doi:10.1186/s12884-020-03124-2
51. Hajizadeh K, Vaezi M, Meedya S, Mohammad-Alizadeh-Charandabi S, Mirghafourvand M. Respectful maternity care and its relationship with childbirth experience in Iranian women: a prospective cohort study. *BMC Pregnancy Childbirth* 2020;20(1):468. doi:10.1186/s12884-020-03118-0
52. Dennis CL, Dowswell T. Psychosocial and psychological interventions for preventing postpartum depression. *Cochrane Database Syst Rev* 2013;2013(2): CD001134. doi:10.1002/14651858.CD001134.pub3
53. Motrico E, Bina R, Kassianos AP, Le HN, Mateus V, Oztekin D, et al. Effectiveness of interventions to prevent perinatal depression: an umbrella review of systematic reviews and meta-analysis. *Gen Hosp Psychiatry* 2023;82:47-61. doi:10.1016/j.genhosppsych.2023.03.007
54. Sangsawang B, Deoisres W, Hengudomsub P, Sangsawang N. Effectiveness of psychosocial support provided by midwives and family on preventing postpartum depression among first-time adolescent mothers at 3-month follow-up: a randomised controlled trial. *J Clin Nurs* 2022;31(5-6):689-702. doi:10.1111/jocn.15928
55. Missler M, van Straten A, Denissen J, Donker T, Beijers R. Effectiveness of a psycho-educational intervention for expecting parents to prevent postpartum parenting stress, depression and anxiety: a randomized controlled trial. *BMC Pregnancy Childbirth* 2020;20(1):658. doi:10.1186/s12884-020-03341-9
56. Fenwick J, Toohill J, Gamble J, Creedy DK, Buist A, Turkstra E, et al. Effects of a midwife psycho-education intervention to reduce childbirth fear on women's birth outcomes and postpartum psychological wellbeing. *BMC Pregnancy Childbirth* 2015;15:284. doi:10.1186/s12884-015-0721-y
57. Marks MN, Siddle K, Warwick C. Can we prevent postnatal depression? A randomized controlled trial to assess the effect of continuity of midwifery care on rates of postnatal depression in high-risk women. *J Matern Fetal Neonatal Med* 2003;13(2):119-27. doi:10.1080/jmf.13.2.119.127
58. Tickle N, Gamble J, Creedy DK. Clinical outcomes for women who had continuity of care experiences with midwifery students. *Women Birth* 2022;35(2):184-92. doi:10.1016/j.wombi.2021.04.004
59. Möller L, Josefsson A, Lilliecreutz C, Gunnervik C, Bladh M, Sydsjö G. Reproduction, fear of childbirth and obstetric outcomes in women treated for fear of childbirth in their first pregnancy: a historical cohort. *Acta Obstet Gynecol Scand* 2019;98(3):374-81. doi: <https://doi.org/10.1111/aogs.13503>