

Systematic Review



A systematic review and meta-analysis of the prevalence of post-traumatic stress disorder (PTSD) in road traffic accident survivors

Kavous Shahsavarinia^{1,2*}, Zahra Sabahi^{1*}, Fateme Tahmasbi¹, Nooshin Milanchian³, Haleh Farshi¹, Javad Mahmoudi⁴, Mostafa Farahbakhsh⁵, Hanieh Salehi-Pourmehr^{1*}, Sakineh Hajebrahimi¹, Homayoun Sadeghi-Bazargani^{5*}

¹Research Center for Evidence-Based Medicine, Iranian EBM Centre: A JBI Centre of Excellence, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

²Emergency and Trauma Care Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

³Department of Internal Medicine, Faculty of Medicine, Tabriz Medical Sciences, Islamic Azad University, Tabriz, Iran

⁴Neurosciences Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

⁵Road Traffic Injury Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

ARTICLE INFO

Article History:

Received: September 12, 2024

Revised: February 09, 2025

Accepted: February 21, 2025

ePublished: November 4, 2025

Keywords:

Accidents traffic, Crashes traffic, Collision traffic, Motor vehicles, Post-traumatic neuroses, Stress disorders, Post-traumatic

*Corresponding Authors:

Hanieh Salehi-Pourmehr,
Emails: salehiha@tbzmed.ac.ir;
poormehr@yahoo.com,
Homayoun Sadeghi-Bazargani,
Email: homayoun.sadeghi@gmail.com

Joint first-authors: Kavous Shahsavarinia and Zahra Sabahi.

Abstract

Background: This systematic review and meta-analysis aimed to evaluate the prevalence of post-traumatic stress disorder (PTSD) among road traffic accident (RTA) survivors, a demographic impacted by over 50 million disabilities globally each year.

Methods: An initial systematic search was conducted in November 2021, with an updated search performed in October 2024. Relevant databases were comprehensively searched using keywords related to "traffic accidents," "road accidents," "motor vehicle accidents," "PTSD," and "Post-Traumatic Stress Disorder." This systematic review and meta-analysis were conducted in accordance with the PRISMA 2020 guidelines and the PICO framework established by Cochrane. The review included studies that involved RTA survivors diagnosed with PTSD, focusing on time frames from one to six months post-accident and utilizing DSM criteria. Eligible studies were reviewed for quality using the standardized critical appraisal instruments from the Joanna Briggs Institute (JBI) Checklist, by two independent reviewers. Meta-analysis was performed using Comprehensive Meta-Analysis statistical software and STATA16 to estimate overall prevalence rates and subgroup analyses to explore variations.

Results: A comprehensive search across multiple databases identified 11,142 articles, of which 92 were reviewed, and 82 were included in the meta-analysis. The findings revealed an overall PTSD prevalence of 20.3% (95% confidence interval [CI]: 18.1%-22.8%; I²:93.86%); 18.7% (95% CI: 16.0%-21.8%; I²: 93.47%) based on clinician-administered assessments and 22.8% (95% CI: 18.8%-27.3%; I²: 93.92%) from self-reported questionnaires. After removing outliers, the total prevalence was decreased to 18.1% (95% CI: 15.4%-21.0%; I²:93.09%), in clinician-administered and 20.8% (95% CI: 17.5%-24.4%; I²: 91.51%) in self-reported questionnaires. Notably, the prevalence was 29.4% (95% CI: 22.4%-37.5%) one-month post-RTA, decreasing to 18.8% (95% CI: 14.8%-23.5%); $P < 0.001$ at three months. Age did not significantly predict PTSD prevalence rates. The quality assessment of the studies included demonstrated moderate to high quality according to the Joanna Briggs Institute standards, ensuring the reliability of the findings. Geographic variability in PTSD prevalence was observed, with lower rates reported in Switzerland, Australia, Germany, and Japan, while higher rates were found in Spain, China, and Iran.

Conclusion: This review highlights a significant PTSD prevalence of 20.3% among traffic accident survivors, emphasizing the need for early intervention and targeted mental health support to mitigate long-term psychological impacts and improve recovery outcomes in this vulnerable population. Screening and public awareness of disease symptoms are recommended.

Introduction

Even with worldwide and local initiatives aimed at preserving life on the roads, approximately 1.35 million people die each year due to road accidents.¹ Road traffic accidents (RTAs) remain the third leading cause of

disability, involving more than 50 million people injured worldwide annually. RTAs, as the most frequent accidental traumatic events, can happen to anyone, including children and teenagers. Evidence has shown that RTAs may increase a person's risk of developing a wide range

of psychiatric disorders, including post-traumatic stress disorder (PTSD), depression, and anxiety.²⁻⁴

PTSD is a chronic and debilitating mental condition that may develop in response to catastrophic life events following exposure to an unintended traumatic event. Over the past few decades, there has been an increase in the number of people affected by traffic accidents worldwide. PTSD is primarily caused by motor vehicle accidents (MVAs) in the general population.⁵ In the United States, approximately six million motor vehicle accidents occur annually, causing over 2.5 million injuries. A study by the National Institute of Mental Health found that more than 39% of those who survive these accidents develop PTSD.⁶ Numerous studies on traumatic events have consistently demonstrated that PTSD can have long-term negative impacts on quality of life. The prevalence of PTSD varies widely between studies, ranging from 4.9% to 34.5%,⁷⁻⁹ which may be related to differences in the time interval between the trauma's occurrence and the assessment of PTSD.

The Diagnostic and Statistical Manual of Mental Disorders-V, fifth edition, outlines that PTSD is characterized by infiltration, avoidance, heightened arousal, and detrimental alterations in mood and cognition.^{10,11} Furthermore, PTSD can result in financial difficulties in addition to physical and psychological harm.¹² Children and adolescents who have experienced RTA often suffer from diminished health-related quality of life due to early signs of PTSD.¹³ Additionally, individuals with PTSD are at an increased risk of metabolic syndrome and obesity.¹⁴ The medical costs for RTA survivors with PTSD are significantly higher than for those without PTSD. Accurate estimation of PTSD prevalence among RTA survivors helps health service providers provide prompt and effective intervention strategies.¹⁵

Several factors are considered potential predictors of PTSD among survivors of MVAs. Previous studies indicate that female sex, depression, a history of RTA, peritraumatic dissociative experiences, an acute stress disorder (ASD) diagnosis, rumination, higher injury severity, and involvement in litigation or compensation following trauma are significant predictors of PTSD.¹⁶⁻¹⁸ Additionally, high levels of emotion (such as fear, helplessness, panic, guilt, or shame) during or right after a traumatic event, a lack of social support following a traumatic event, and previous psychological adjustment issues are all factors that increase the risk of developing PTSD.¹⁷ While injuries to the driver or passengers did not receive much support as predictive factors, the individual's perceptions and responses to the accident, avoidance behavior, and suppressed thoughts about the accident significantly predicted PTSD.¹⁶

A comprehensive understanding of PTSD prevalence is essential for the development of effective treatment strategies, informing policy decisions, and ultimately enhancing mental health outcomes for both individuals and communities. Two prior systematic reviews have

been published to assess the prevalence of PTSD after RTA and to identify the predictors of PTSD in adult survivors of road traffic collisions.^{15,19} This systematic review and meta-analysis aim to investigate the prevalence of PTSD among survivors of RTA by updating previously published systematic reviews.

Methods

This systematic review and meta-analysis was conducted in accordance with the PRISMA 2020 guidelines, which provide a standardized framework for transparent and comprehensive reporting of systematic reviews and meta-analyses.²⁰ The PRISMA checklist was followed to ensure that all critical elements of the review process were addressed, including the identification, screening, eligibility, and inclusion of studies.

Study Population

Following the PICO framework established by Cochrane, we have previously articulated our intention to conduct a systematic review focusing on survivors of RTAs (Population) to investigate and synthesize the total prevalence (Outcome).

Inclusion/Exclusion Criteria

The following criteria were considered for eligibility: the sample consisted of RTA survivors diagnosed with PTSD 1 to 6 months after the accident (using different versions of DSM criteria or self-reported questionnaires). Studies were excluded if they were presented in non-English, were conference abstracts, were case studies or dissertations, were letters, or reviews. Additionally, while a month or more should pass before a PTSD assessment is conducted (no more than six months in line with DSM-IV criteria), studies on delayed PTSD and cases influenced by confounding variables, such as traumatic brain injury or post-traumatic amnesia, were excluded. No age restrictions were considered.

Information Sources

An initial systematic search was conducted in November 2021, with an updated search performed in October 2024 in PubMed, Ovid, ProQuest, Scopus, Web of Science, Cochrane Library, and Google Scholar using the keywords Traffic Accident, Traffic Collision, Traffic Crash, Road accident, Motorcycle accident, Motorcar accident, Motor vehicle accident, PTSD, Moral Injury, and Post-traumatic Neuroses. The updated systematic search was conducted in October 2024 using the same search strategy and sources. Also, the reference lists of the studies were checked to retrieve any relevant publications. The full version of the search strategy is presented in [Supplementary file 1](#).

Study Selection

First, all identified citations were loaded into EndNote X20. After deleting duplicates, two independent reviewers

screened titles and abstracts to assess the review's inclusion criteria. Then, two independent professional reviewers (HS and ZS) evaluated the full text of the selected eligible studies in detail. If any did not meet the inclusion criteria, they were excluded. Any reviewer disagreements were resolved through discussion or by a third reviewer (HSB).

Data Extraction

Utilizing the modified standard JBI data extraction tool, two reviewers (HS and ZS) independently determined the qualified papers and retrieved their data. Any disagreements were settled by consensus between the two reviewers or by conversing with the third reviewer (HSB). The data extraction table includes the study's first author, publication year, the study's nation, sample size, the timing of the PTSD assessment, the seriousness of the injury, the PTSD assessment tool, and prevalence. The PTSD diagnosis was determined through clinician assessments or self-reported questionnaires. The clinician-based evaluation contributed to the PTSD diagnosis, while the self-reported questionnaires indicated probable PTSD diagnosis.

Assessment of Methodological Quality

According to standardized critical appraisal instruments from the Joanna Briggs Institute, eligible studies were critically appraised by two independent reviewers (HS and ZS) at the study level. Any disagreements were resolved by discussion or consultation with the third reviewer (HSB). Studies with a half or higher score in questions were included as high or moderate-quality studies (available at: <https://jbi.global/critical-appraisal-tools>).

Statistical Analysis

All analyses were performed using registered copies of Comprehensive Meta-Analysis statistical software (version 3; Biostat, Englewood, NJ) and STATA16 (StataCorp, College Station, TX, USA). The study statistician extracted the data for the primary outcomes. The random effect model was employed because we might not have access to other unidentified, unregistered, or unpublished investigations. The between-study heterogeneity was assessed using statistics such as the Cochran Q test, Tau-squared, and I-squared. Significant results of the test and values higher than 75% for I-squared were considered substantial heterogeneity.²¹ Estimates were made for the effect sizes and 95% CIs. The publication bias was evaluated using funnel plots. To assess the bias, Egger's²² and Begg's²³ as well as Duval and Tweedie's trim and fill were performed. The age of the research population was used as the independent variable in the meta-regression analysis, which is typically necessary to identify the cause of heterogeneity.

As needed, subgroup analysis based on the PTSD assessment tool by country, age groups (adults > 18 years, children < 18 years),²⁴ and gender was conducted to identify the sources of heterogeneity.

Results

Study Inclusion

Our initial search retrieved 11142 articles from databases. After removing duplicates (n = 5121), reviewing the titles/abstracts, and reading the full text of eligible articles, 598 full texts of the articles were evaluated. Finally, 92 studies were systematically reviewed,^{3,4,7,16,25-112} and 82 studies (including 30 clinician-administered measures studies) were candidates for meta-analysis (Figure 1).

Characteristics of Included Studies

The full details of the included studies are presented in Table 1. The prevalence of PTSD was evaluated in 82 studies.

The research encompassed a diverse range of countries, including Saudi Arabia,²⁵ Albania,⁹³ Australia,^{32,42,49,54,66} Belgium,⁸⁶ China,^{51,62,69,79,103} Scotland,⁹² Denmark,⁴⁶ Ethiopia,^{28,29,41,84} France,^{16,35,77,81,95,111} Germany,^{7,26,31,45,60} Greece,^{98,106} India,^{73,110} Iran,^{57,76} Taiwan,⁸² Israel,^{61,67} Japan,^{4,63,70,105} Malaysia,²⁷ Nigeria,⁵³ Portugal,⁷⁴ South Korea,⁸⁷ South Africa,^{71,78} Spain,³⁶ Sweden,³⁸ Switzerland,¹⁰⁴ Turkey,⁸³ the United Kingdom,^{3,39,40,50,55,64,65,68,90,96,97,99,101} Balkan,⁵⁹ Greece, Germany, and Italy,⁷² and the United States.^{30,34,37,43,44,47,48,52,56,58,75,80,100,102,107,108,109,112} Some studies reported the prevalence of PTSD in children.^{7,85,90-92,93-106}

The studies varied in geographical distribution, with 39 conducted in Europe, 20 in Asia, 19 in North America, 7 in Oceania, and 7 in Africa. Sample sizes were notably heterogeneous, ranging from a minimum of 21^{77,93} to a maximum of 3,604 participants,¹⁰² and assessment durations spanned from several days to six months. The severity of injuries was documented through qualitative measures.

Diagnostic assessments predominantly utilized standardized instruments such as the Clinician-Administered PTSD Scale (CAPS) aligned with DSM-IV criteria, the PTSD Checklist-Civilian version (PCL-C), the Structured Clinical Interview for DSM (SCID) as per DSM-III-R or DSM-IV standards, the Impact of Event Scale-Revised (IES-R), the PCL-S, the Composite International Diagnostic Interview for PTSD (CIDI-PTSD), the PTSD Diagnostic Scale (PDS), the Perceived Stress Scale (PSS), the Penn Inventory, and the PTSD of the Mini International Neuropsychiatric Interview (M.I.N.I). Although less commonly employed, additional evaluation methods are comprehensively outlined in Table 1.

Among the studies reviewed, PTSD diagnoses were self-reported in 35 cases and clinician-administered in 50 cases. The reported prevalence of PTSD varied widely, falling within the range of 2.9% to 77.8%. Furthermore, the mean age of participants spanned from 23 years to 66.09 years, with a standard deviation of 5.9 years. The time point to measure PTSD was between 1 to 6 months after RTAs in survivors.

We analyzed the frequency of PTSD according to the clinician-administered or self-reported questionnaire in




Figure 1. The PRISMA flow diagram shows the search and screening processes

the included studies at different time points. According to the results of clinician-administered assessment, the prevalence of PTSD varied between the minimum percent of 13.3% in Ongecha-Owuor et al⁷¹ to 48% at the Lesaca et al⁸⁵ study one month after RTA. At six weeks after RTA, the prevalence of PTSD varied between 10.4% in the Irish et al study⁵² to 37.1% in the Stallard study. The minimum and maximum rate was 7.1%¹⁰⁴ to 51%⁸¹ at two months post-injury, respectively. At three months post-injury, this rate was 6.8%,⁶⁷ and 50%,⁶¹ and at 6 months 1.7%,⁹⁷ and 38.9%,⁷⁷ respectively.

In terms of self-administered questionnaires, the prevalence rate of PTSD was low in the study of Allenou et al from France (3.6%)⁹⁵ at one month post-injury, while Fekadu et al⁴¹ reported the highest rate (46.5%). At 6 weeks post-injury, Angerpointner et al²⁶ reported 5.6% of PTSD, and Wang et al⁸² reported 82.8% of cases with this disorder. At three months, the Angerpointner et al²⁶ study found that only 2.8% of cases had PTSD, while Bryant et al⁹⁰ from the UK, and Schäfer et al⁷ from Germany reported 25% of its prevalence.

The minimum rate of PTSD after RTA was observed in the Mehnert et al⁸⁹ study (8.0%), and the maximum rate was in the Papadakaki et al⁷² study (39.6%).

Clinician-Administered Measures

We calculated the total prevalence of PTSD according to the clinician-administered measures, which was 18.7% (95% CI: 16.0%-21.8%; I²: 93.47%) (Figure 2).

After removing outliers, the total prevalence decreased to 18.1% (95% CI: 15.4%-21.0%; I²: 93.09%) in clinician-administered.

In subgroup analysis based on the time points in clinician-administered measures-related studies, the results revealed that 1 month after RTA, the prevalence of PTSD was 29.4% (95% CI: 22.4%-37.5%; I²: 85.97%) among 11 eligible studies; in 3 months following RTA, this rate was 18.8% (95% CI: 14.8%-23.5%; I²: 89.83%) in 13 included studies, and at six months was 13.0% (95% CI: 9.6%-17.3%; I²: 95.81%) in the 24 included studies. At the other time point, consisting of 2 months following RTA, this rate was 26.5% (95% CI: 11.2%-50.5%; I²: 88.52%) in 4 eligible studies, and at 4 months was 36.6% (95% CI: 30.6%-43.0%; I²: 11.49%) in two eligible studies (Figure 3).

In subgroup analysis based on continents in clinician-administered measures-related studies, the prevalence in American areas was 18.3% (95% CI: 13.3%-24.7%) among 14 studies; in European regions, 19.5% (95% CI: 15.5%-24.1%) in 19 eligible studies, 18.2% (95% CI: 12.7%-25.4%) among six related studies in Western Pacific regions, 19.9% in Asia (95% CI: 12.8%-29.6%) in nine eligible studies, and 14.8% (95% CI: 11.12%-19.3%) in 3 African studies (Figure 4).

According to the subgroup analysis based on the different countries within the clinician-administered group, the most published papers were from the USA (n=14 studies, with a prevalence of 23.8% (95% CI: 16.2%-33.6.2%; I²: 98.8%). According to our findings, the

Table 1. The characteristics of included studies

First author (year)	Country	Sample size	Timepoint	Injury severity	Measure to assess PTSD and DSM version	Self-reported/ Clinician administered	PTSD prevalence	Mean age
Alshardan (2020) ²⁵	Saudi Arabia	334	NM	NM	PCL-C (DSM-IV)	Self-reported	39.2	
Angerpointner (2020) ²⁶	Germany	36	6 weeks 3 months	Minor	IES-R	Self-reported	5.6 2.8	39.8 (16.0)
Bahari (2017) ²⁷	Malaysia	68	1 month	Minor, moderate, and major	Malay Post-Traumatic Stress Disorder Checklist Civilian version	Self-reported	7.4 (incidence)	66.09 (5.9)
Bedaso (2020) ²⁸	Ethiopia	416	Not less than one month	Hospitalized	PCL-S	Self-reported	15.4	
Berna (2012) ¹⁶	France	155	6 months	Hospitalized	CAPS (DSM-IV)	Clinician administered	7.74	36.7 (16.4)
Bezabh (2018) ²⁹	Ethiopia	603			PCL-C	Self-reported	All emergency responders: 19.9 (ambulance nurses:11.5; firefighters:20.7)	
Blanchard (1995) ³⁰	USA	158	4 months	Sought medical attention	SCID (DSM-III-R)	Clinician administered	39.2	MVA victims: 35.4 (12.5) control: 37.7 (14.00)
Brand (2014) ³¹	Germany	258	NM	Sought medical attention	Criterion A: Exposed to: death, threatened death, serious injury, and sexual violence (DSM-V)		0.78 (incidence)	
Bryant (2003) ³²	Australia	87	6 months	Hospitalized	CIDI-PTSD (DSM-III-R)	Clinician administered	22	Male: 29.95 (11.49) female:33.36 (13.28)
Bryant (2000) ³³	Australia	113	6 months	Hospitalized >24 h	CIDI-PTSD (DSM-III-R)	Clinician administered	21	No-TBI: 33.70 (11.98)
Buckley (2004) ³⁴	USA	65	1 month	serious	SCID (DSM-IV)	Clinician administered	17	36.05 (15.04)
Chossegros (2011) ³⁵	France	541	6 months	Hospitalized	PCL (DSM-IV)	Self-reported	18	
Coronas (2011) ³⁶	Spain	119 108	1 month 4 months	Serious	SCID (DSM-III-R)	Clinician administered	45.4 32.8	38.3 (12.3)
Delahanty (2003) ³⁷	USA	59	1 month	Serious	SCID (DSM-IV)	Clinician administered	20	37.3 (17.7)
Doohan (2017) ³⁸ (bus)	Sweden	54	1-3 months	Minor, moderate, severe	TSQ		31% were high risk for PTSD	57
Ehlers (1998) ³⁹	UK	888	3 months	None, mild & moderate	PSS (DSM-IV)	Clinician administered	23.1	33.4 (13.1)
Ehring (2008) ⁴⁰	UK	141	6 months	Moderate to severe	SCID (DSM-IV)	Clinician administered	12.1	34.95 (10.60)
Fekadu (2019) ⁴¹	Ethiopia	299	1 month	No major trauma	PCL-C	Self-reported	46.5	The median age was 31 with (IQR) of 25–42.
Fitzharris (2006) ⁴²	Australia	62 Males = 35 Females = 27	6-8 weeks 6-8 months	Hospitalized	PCL-C	Self-reported	Male: 2.9%, Female:7.4% Male: Nil, Female:7.4%	males: 35.3 (12.3), Females: 38.7 (12.3)
Flesher (2001) ⁴³	USA	70	1 month	Hospitalized	SCID (DSM-IV)	Clinician administered	17	33.2 (14.6)
Fredman (2017) ⁴⁴	USA	114	4 weeks 16 weeks	Severe	PCL-C	Self-reported	42.1 24.3	38.14 (12.52)
Frommberger (1998) ⁴⁵	Germany	152	6 months	Hospitalized minimum of bone fracture	IES, PSS (DSM-III-R)	Self-reported	18.4	
Fuglsang (2004) ⁴⁶	Denmark	90	6–8 months	Attended ED	PDS (DSM-IV)	Self-reported	17	33.99 (11.3)
Fullerton (2001) ⁴⁷	USA	122	1 month	Serious	SCID (DSM-III-R)	Clinician administered	34.4	35.6 (13.1)

Table 1. Continued.

First author (year)	Country	Sample size	Timepoint	Injury severity	Measure to assess PTSD and DSM version	Self-reported/ Clinician administered	PTSD prevalence	Mean age
Gabert-Quillen (2012) ⁴⁸	USA	201	6 months	hospitalized	CAPS	Clinician administered	7	39.6 (15.7)
Hamanaka (2006) ⁴	Japan	82	6 months	Serious	SCID (DSM-IV)	Clinician administered	8.5	
Harvey (1998) ⁴⁹	Australia	71	6 months	Hospitalized > 24 h	CIDI-PTSD (DSM-III-R)	Clinician administered	25.4	33.29 (12.00)
Holeva (2001) ⁵⁰	UK	265	4–6 months	Serious	Penn Inventory (DSM edition not specified)	Self-reported	23	
Hu (2018) ⁵¹	China	70	6 months	mild	CAPS (DSM-IV)	Clinician administered	41.4	
Irish (2011) ⁵²	USA	196	6 weeks	Hospitalized	CAPS (DSM-IV)	Clinician administered	10.36	38.4 (14.7)
			6 months	Mild, moderate & severe			7.14	
Iteke (2011) ⁵³	Nigeria	150	1-12 months		PTSD module of the Mini International Neuropsychiatric Interview (M.I.N.I.)	Clinician administered	26.7	RTA:31.61 ± 9.18, control 1: 32.14 ± 8.85, control 2: 33.01 ± 8.95
Jeavons (2000) ⁵⁴	Australia	72	3 months	Attended to hospital	PTSD-I (DSM-III-R)	Clinician administered	8.3	31.8 (12.78)
		62	6 months				8	
Jones (2005) ⁵⁵	UK	131	6 weeks	Severe	PSS (DSM-IV)	Clinician administered	Non-TBI: 27.4 TBI: 30.4	36.75 (12.77)
			3 months				Non-TBI: 18 TBI: 17.2	
Kassam-Adams (2009) ⁵⁶	USA	251 parents of children with RTA	6 months	Hospitalized	PTSD Checklist		8	
Khodadadi-Hassankiadeh (2017) ⁵⁷	Iran	528	6 weeks-6 months	Attended to hospital	PSS	Clinician administered	30.49	33.59 (13.29)
Kobayashi (2019) ⁵⁸	USA	280 (120 women, 160 men)	6 weeks	Admitted to level-1 trauma centers	CAPS (DSM-IV)	Clinician administered	Women: 19.2 Men: 8.1	women:39.93 (15.29), men : 37.41 (15.00)
		217 (99 women, 118 men)	6 months				Women: 14.1 Men: 2.5	
Kovacevic (2021) ⁵⁹	Balkans	200	1 month	Mild, moderate, serious, severe, critical	PCL-C	Self-reported	35.5	
			6 months				20.5	
Kuhn (2006) ⁶⁰	Germany	58	6 months	Moderate to Severe	SCID (DSM-IV German version)	Clinician administered	6	38.6 (13.6)
Kupchik (2007) ⁶¹	Israel	60	3 months	Outpatient clinic	CAPS-2, SCID-I/P	Clinician administered	50	PTSD: 44.6 (11.1) non-PTSD: 45.4 (13.2)
Li (2021) ⁶²	China	206	4–12 months	Mild, severe, critically severe	PCL-S	Self-reported	24.8	39.8 (12.5)
Matsuoka (2008) ⁶³	Japan	100	4–6 weeks	Severe	CAPS (DSM-IV)	Clinician administered	8	37.0 (16.1)
Mayou (1993) ⁶⁴	UK	174	3 months	Minor & Major	Diagnostic criteria for PTSD (DSM-III-R)	Clinician administered	8	
Mayou (1997) ⁶⁵	UK	111	3 months	Attended ED		Clinician administered	10	
McFarlane (1997) ⁶⁶	Australia	26	6 months	Hospitalized	CAPS (DSM-IV)	Clinician administered	26.9	
Naim (2014) ⁶⁷	Israel	415	3 months	Minor, admitted to ED	CAPS, PCL, CADSS	Clinician administered	6.75	
Nightingale (2000) ⁶⁸	UK	60	6 weeks		PDS (DSM-IV)	Self-reported	30.8	T1:33.3 (10.5), T2:34.8 (10.8)
Ning (2017) ⁶⁹	China	166	3 months		PCL-C	Self-reported	15	38.75 (1.13)
Nishi (2013) ⁷⁰	Japan	106	6 months	Admitted to ICU	CAPS	Clinician administered	7.5	38.3 (16.0)

Table 1. Continued.

First author (year)	Country	Sample size	Timepoint	Injury severity	Measure to assess PTSD and DSM version	Self-reported/ Clinician administered	PTSD prevalence	Mean age
Ongecha-Owuor (2004) ⁷¹	Africa	264	1 month	Serious	SPI (DSM-IV)	Clinician administered	13.3	
Papadakaki (2017) ⁷²	Greece, Germany and Italy	initial:120 (Greece=41, Germany=3, Italy=40), 12 months: 93	6, 12 months	Admitted to ICU	IES-R	Self-reported	PTS: Baseline: 43.5% 6 months: 39.6% 12 months: 21.1%	41.8 (16.7)
Prakasam (2013) ⁷³	India	86	6 months	Moderate, major	IES-R	Self-reported	23.3	
Pires (2013) ⁷⁴	Portugal	124	4 months	serious	RTES		58.90	
Ryb (2009) ⁷⁵	USA	367	6 months	Hospitalized	Diagnostic criteria for PTSD (DSM-IV)	Clinician administered	27.5	
Saberi (2013) ⁷⁶	Iran	385	NM		PCL-C Persian version(	Self-reported	19.2	35.45±9.04
Shaikh (2012) ⁷⁷	France	21	2 months	hospitalized	CAPS	Clinician administered	33.3	At 2 months PTSD + : 23, PTSD - :26
		18	6 months				38.9	at 6 months PTSD + : 29, PTSD - :20
Smith (2007) ³	UK	39	4 months	Minor (out-patients)	SRS-PTSD (DSM-III-R), IES	Self-reported	12.8	
Suliman (2014) ⁷⁸	South Africa	Initial:131, 3 months: 104, 6 months: 101	3 months, 6 months	Minor, major	CAPS	Clinician administered	baseline: 22.9% 3 months: 19.6% 6 months: 12.2%	PTSD: 34.75 (11.54) no PTSD: 33.71 (11.16)
Sun (2013) ⁷⁹	China	62	6 months		CAPS	Clinician administered	33.9	Trauma-exposed victims with PTSD (N=21): 40.86±12.26 Trauma-exposed victims without PTSD (N=17): 35.64±11.91 Follow-up of trauma exposed victims with PTSD (N=11): 42.09±12.79 Healthy control: 40.23±12.54
Ursano (1999) ⁸⁰	USA	122	1 month	Serious	SCID (DSM-III-R)	Clinician administered	34.4	35.6 (13.1)
		99	3 months				25.2	MVA: 35.59 (13.06), control
		99	6 months				18.2	37.16 (13.09)
Vaiva (2003) ⁸¹	France	123	2 months	Hospitalized	CAPS (DSM-IV)	Clinician administered	51	
Wang (2005) ⁸²	Taiwan	64	1 week	Hospitalized	PTSD-RI (DSM-III-R)	Self-reported	87.5	33 (11.77)
			6 weeks				82.8	
Yasan (2009) ⁸³	Turkey	84	3 months	Attended ED	CAPS (DSM-IV)	Clinician administered	29.8	
		78	6 months				23.1	
Yohannes (2018) ⁸⁴	Ethiopia	492	1 month		PCL-S	Self-reported	22.8	30.12 (7.02)
Children								
Bryant (2004) ⁹⁰	UK	86	3 months	Minor, or hospitalized	Post-Traumatic Stress Disorder Reaction Index (RI) (DSM-IV)	Self-reported	25	12.27 (2.86)
			6 months				18	
DI Gallo (1997) ⁹¹	Scotland	49	12-15 weeks		PTSD-RI; IES	Self-reported	Mild: 35; moderate: 8; severe: 6	10.2 (3.6)

Table 1. Continued.

First author (year)	Country	Sample size	Timepoint	Injury severity	Measure to assess PTSD and DSM version	Self-reported/ Clinician administered	PTSD prevalence	Mean age
Gillies (2003) ⁹²	Scotland	158	baseline	Attended ED	CPTS-RI, CAPS-C	Clinician administered	Mild: 48 Moderate: 18 Sever: 0	
			2-16 days				Mild: 33 Moderate: 7 Sever: 7	
			12-15 weeks				Mild: 44 Moderate: 22 Sever: 7	
Jones-Alexander (2005) ⁹³	Albany	21			CPTSDI, PCL-C		38.1	12.7
Landolt (2005) ⁹⁴ (PTSS)		68	4–6 weeks	Hospitalized	Child PTSD Reaction Index (RI)(DSM-IV-TR); PDS for parents		16.2; mothers (20%); fathers (11.3%)	Children: 9.82 (2.55)
Allenou (2010) ⁹⁵	France	Mothers: 72 Fathers: 28	5 weeks	NM	PCL-S	Self-reported	18.1 in mothers 3.6 in fathers	Age of fathers: 40.9 (5.3); Age of mothers 41.7 (6.2)
Meiser-Stedman (2009) ⁹⁶	UK	28	2-4 weeks 6 months	Attended ED	RIES-C; CPTC	Self-reported	21.4 10.7	13.2 (1.9)
Mirza (1998) ⁹⁷	UK	119	6 months	Attended ED	FRI and the PTSD symptom checklist (DSM-IV).	Clinician administered	Severe: 12 moderate:1.7 mild:3.4	13.61 (2.44 years)
Pervanidou (2007) ⁹⁸	Greece	56	1 month	minor, moderate and serious	K-SADS; CPTS-RI	Clinician administered	41.1 Boys: 32 Girls: 7	children: 10.70 (2.46), control: 10.49 2.59
		48	6 months				18.8 Boys: 16 Girls: 2	
Salter (2004) ⁹⁹	UK	67	few weeks	Admitted to hospital	CAPS-C	Clinician administered	37	15 (3)
Schäfer (2006) ⁷	Germany	72	1 week	Attended ED	Impact of Event Scale – Revised (IES-R)	Self-reported	11	13.6(3.3)
		69	3 months				25	
Stallard (2001) ¹⁰⁰	USA	97	6 weeks	Attended ED	CAPS-C	Clinician administered	37.1	14.62 (3.16)
Stallard (2004) ¹⁰¹	UK	158	4 weeks	Attended ED	CAPS-C	Clinician administered	29.1	14.85 (3.11)
Williams (2015) ¹⁰²	USA	3604	6 months	Serious	NWS	Clinician administered	7.4	14.63 (1.66)
Wu (2016) ¹⁰³	China	537	3 months	Admitted to the hospital	CAPS-CA	Clinician administered	24.77	6.8±0.9
Zehnder (2010) ¹⁰⁴	Switzerland	50	2 months	Hospitalized	CAPS-CA	Clinician administered	7.1	7-16 years
		50	6 months				4	
Maeda (2009) ¹⁰⁵ (ship)	Japan	Adolescent: 9 Adult: 17	2 months		CAPS	Clinician administered	Adolescent: 77.8 Adult: 12	student: 17.0 (0.0), crew: 45.9 (11.6)
Giannopoulou (2021) ¹⁰⁶	Greece	168	2 months		Children's Revised Impact of Events Scale (CRIES-13)	Self-reported	78	14.5 (1.3)
Ziobrowski (2021) ¹⁰⁷	USA	1003	3 months		CAPS-DSM-IV PTSD	Clinician administered	26.60	34.5 [24-43]
Kessler (2021) ¹⁰⁸	USA	666	2 months		PCL-5	Self-reported	39.00	-
Joormann (2022) ¹⁰⁹	USA	1306	3 months		PCL-5	Self-reported	20	-
Arora (2021) ¹¹⁰	India	250	1-12 months		PCL-5	Self-reported	32.40	31-45
Yrondi (2022) ¹¹¹	France	125	5 weeks, 6 months		PCL-5	Self-reported	5 weeks: 13.6, 6 months: 10.3	40.83 (5.21)
Neylan (2021) ¹¹²	USA	666	2 months		PCL-5	Self-reported	42	-

Airplane

Table 1. Continued.

First author (year)	Country	Sample size	Timepoint	Injury severity	Measure to assess PTSD and DSM version	Self-reported/ Clinician administered	PTSD prevalence	Mean age
Lesaca (1996) ⁸⁵	USA	Trauma counseling: 21 no trauma counseling:20	4 weeks	NM	DSM-IV	Clinician administered	48 10	
		Trauma counseling: 21 no trauma counseling: 20	8 weeks				24 0	
		No trauma counseling: 20	12 weeks				14 25	
Train accident								
Engelhard (2002) ⁸⁶	Belgium	Directly exposed: 29	3 weeks 3.5 months	Serious	PSS	Clinician administered	28 17	53 (17)
Kim (2013) ⁸⁷ (subway drivers)	South korea	826	NM	NM	K-CIDI 2.1	Clinician administered	1.5	
Lemos (2018) ⁸⁸	Portugal	216	Baseline		PCL-C (Portuguese version)	Self-reported	8.3	
			Less than a month				37.5	
			6 months				10.2	
Mehnert (2012) ⁸⁹	Germany	71	1 month		Posttraumatic Diagnostic Scale (PDS) — German version	Self-reported	moderate PTS: 28%, moderate to severe PTS:42%, severe PTS :11%	48 (7.8)
		49	6 months				moderate PTS: 29%, moderate to severe PTS :29%, severe PTSD: 8%	

lowest prevalence was observed in Switzerland (5.8%), Germany (6%), and Japan (8%) in one, one, and three eligible studies. In contrast, Spain (39%), China (32.2%), Iran (30.5%), Greece (29.3%), and France (28.9%) had the highest prevalence in two, three, one, two, and four eligible studies, respectively (Figure 5). Furthermore, the distribution of RTA survivors is schematically presented in Figure 6.

In subgroup analysis to show the prevalence of PTSD based on the injury severity within the clinician-administered group, in hospitalized patients, the prevalence rate was 18.7% (95% CI: 12.7%-26.6%); in cases attended to the emergency department, this rate was 16.7% (95% CI: 10.7%-25.2%) and in serious injuries 20.3% (95% CI: 14%-28.7%) (Supplementary file 2, Figure S1).

In subgroup analysis based on age within the clinician-administered group, the prevalence of PTSD in adults (age > 18 years) was 19.2% (95% CI: 16.4%-22.4%; I²: 90.10%), and in children (age < 18 years) was 17.4% (95% CI: 11.9%-24.8%; I²: 96.10) (Figure S2). After removing outlier studies, the prevalence was estimated to be 17.8% (95% CI: 15.2%-20.8%; I²: 88.88%) in adults.

Figure S3 represents the result of the subgroup analysis

based on the checklist used for diagnosis.

Self-Reported Measures

For self-reported PTSD, the prevalence rate was 22.8% (95% CI: 18.8%-27.3%; I²: 93.92%); After removing outliers, the total prevalence was decreased to 20.8% (95% CI: 17.5%-24.4%; I²: 91.51 %) using self-reported questionnaires (Figure 2 and Figure S4).

In subgroup analysis based on the time points in self-reported group studies, 1 month after RTA, the prevalence of PTSD was 22.6% (95% CI: 15.9%-31.2%; I²: 92.61%) among 10 eligible studies; in 3 months following RTA, this rate was 19.7% (95% CI: 15.2%-25%; I²: 58.08) in 5 included studies, and at six months was 17.4% (95% CI: 15.8%-23%; I²: 83.06%) in the 10 included studies. At the other time point, consisting of 2 months following RTA, this rate was 53.4% (95% CI: 36.5%-69.6%; I²: 97.27%) in 3 eligible studies, and at 4 months was 19.6% (95% CI: 15.8%-33.6%; I²: 54.69%) in two eligible studies (Figure S5).

In subgroup analysis based on the WHO regions in self-reported group studies, the prevalence in American areas was 31.6% (95% CI: 29.9%-33.4%) among five studies; in European regions, 24.4% (95% CI: 22.7%-26.1%) in 18




Figure 2. The forest plot for the prevalence of PTSD based on Clinician-administered and self-reported groups




Figure 3. The forest plot for the prevalence of PTSD at different time points within Clinician-administered group




Figure 4. The forest plot for the prevalence of PTSD in different WHO regions within Clinician-administered group




Figure 5. The forest plot for the prevalence of PTSD in different countries within Clinician-administered group

eligible studies, 5.4% (95% CI: 1.7%-15.4%) among two related studies in Western Pacific regions, 25.5% in Asia (95% CI: 21.8%-29.7%) in 5 eligible studies, and 27.5% (95% CI: 24.9%-30.3%) in 3 African studies (Figure S6).

According to the subgroup analysis based on the different countries, the most published papers were from the UK (n=5 studies, with a prevalence of 22.1% (95% CI: 18.1%-26.2%; I^2 : 23.19%). According to our findings, the lowest prevalence was observed in Australia (5.4%), Malaysia (7.4%), and Portugal (10.2%) in eligible studies in each country. In contrast, Taiwan (82%), Greece (78%), and the USA (32.7%) had the highest prevalence in one, one, and four eligible studies, respectively (Figure S7).

We performed a subgroup analysis to show the prevalence of PTSD based on the injury severity. The results showed that in hospitalized patients, prevalence rates of PTSD following an RTA varied considerably across studies, ranging from 9% to 43.1% (point estimate of 21.5%); in cases attended to the emergency department, this rate was 19.7% (95% CI: 14.8%-25.8%). The prevalence rates ranged from 14.8% to 36.8% (point estimate of 24%) in the studies that included severe injury cases (Figure S8).

In subgroup analysis based on age within the self-reported group, the prevalence of PTSD in adults (age > 18 years) was 23.3% (95% CI: 18.8%-28.5%), and in children (age < 18

years) was 21.5% (95% CI: 16.4%-27.6%) (Figure S9).

Combining the Studies with Clinician-Administered Measures and a Self-Reported Checklist for PTSD Assessment

To evaluate the publication bias, a funnel plot was drawn, and according to the results of the Egger's regression test, there was significant publication bias among the included studies ($P < 0.001$) (Figure S10).

Meta-regression models were used to investigate further the contribution of a variable to the prevalence heterogeneity. The results of this multivariate analysis suggested that the mean age of RTA survivors was not associated with significant heterogeneity between prevalence rates of PTSD ($P = 0.711$) (Table 2, Figure S11).

Methodological Quality

Eligible studies were reviewed using the JBI Evidence Quality Evaluation Checklist. The results show that all included studies were of moderate to high quality (more than 60% "yes" response). The detailed results of the quality assessment are presented in Table 3.

Discussion

Our comprehensive meta-analysis, encompassing 82 studies identified from an initial 11,142 articles, revealed an overall pooled prevalence of PTSD of 20.3% (95%




Figure 6. The distribution of RTA survivors on a global scale

Table 2. Multivariate meta-regression of included studies

Meta-regression		Number of observations = 24			
REML estimate of between-study variance		tau2 = 0.01963			
% Residual variation due to heterogeneity		I-squared res: 94.34%			
Proportion of between-study variance explained		Adj R-squared = -1.62%			
ES	exp(b)	SE	t	P > t	[95% Confidence Interval]
Mean age	-0.0008	0.0022	-0.37	0.711	[-0.0052, 0.0035]
cons	0.2276	0.0755	3.01	0.004	[0.0764, 0.3787]

Table 3. Methodological quality assessment using the JBI Evidence Quality Evaluation Checklist (<https://jbi.global/critical-appraisal-tools>)

	Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	Q9	Q10	Q11	Q12	Q13
Cohort studies													
Angerpointner (2020) ²⁶	NA	NA	Yes	Unclear	No	Unclear	Yes	Yes	Yes	Yes	Unclear		
Fitzharris (2006) ⁴²	NA	NA	Yes	Unclear	No	Unclear	Yes	Yes	Yes	Yes	Unclear		
Bryant (2003) ³²	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Ehring (2008) ⁴⁰	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Chossegros (2011) ³⁵	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Li (2021) ⁶²	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Coronas (2011) ³⁶	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Delahanty (2003) ³⁷	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes					
Doohan (2017) ³⁸ (bus)	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		
Ehlers (1998) ³⁹	NA	NA	Yes	Unclear	Unclear	Unclear	Yes	Yes	Yes	Yes	Yes		
Flesher (2001) ⁴³	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes					
Fekadu (2019) ⁴¹	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Kessler (2021) ¹⁰⁸	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Fredman (2017) ⁴⁴	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Frommberger (1998) ⁴⁵	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Fuglsang (2004) ⁴⁶	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Fullerton (2001) ⁴⁷	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Giannopoulou (2021) ¹⁰⁶	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Ziobrowski (2021) ¹⁰⁷	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Hamanaka (2006) ⁴	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Harvey (1998) ⁴⁹	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Holeva (2001) ⁵⁰	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Hu (2018) ⁵¹	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Kovacevic (2021) ⁵⁹	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Jeavons (2000) ⁵⁴	NA	NA	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Jones (2005) ⁵⁵	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Yrondi (2022) ¹¹¹	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Kassam-Adams (2009) ⁵⁶	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Bahari (2017) ²⁷	Yes	Yes	Yes	Unclear	Unclear	Unclear	Yes	Yes	Yes	Yes	Yes	Unclear	
Irish (2011) ⁵²	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Gabert-Quillen (2012) ⁴⁸	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Kobayashi (2019) ⁵⁸	NA	NA	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Kuhn (2006) ⁶⁰	Yes	Yes	Yes	Unclear	Unclear	Unclear	Yes	Yes	Yes	Yes	Yes	Unclear	
Schäfer (2006) ⁷	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Yasan (2009) ⁸³	NA	NA	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Allenou (2010) ⁹⁵	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Mayou (1993) ⁶⁴	NA	NA	Yes	Unclear	Unclear	Unclear	Yes	Yes	Yes	Yes	Yes	Unclear	
Bryant (2000) ³³	Yes	Yes	Yes	Unclear	Unclear	Unclear	Yes	Yes	Yes	Yes	Yes	Unclear	
Bryant (2004) ⁹⁰	NA	NA	Yes	Unclear	Unclear	Unclear	Yes	Yes	Yes	Yes	Yes	Unclear	
Buckley (2004) ³⁴	NA	NA	Yes	Unclear	Unclear	Unclear	Yes	Yes	Yes	Yes	Yes	Unclear	
DI Gallo (1997) ⁹¹	NA	NA	Yes	Unclear	Unclear	Unclear	Yes	Yes	Yes	Yes	Yes	Unclear	
Gillies (2003) ⁹²	NA	NA	Yes	Unclear	Unclear	Unclear	Yes	Yes	Yes	Yes	Yes	Unclear	
Brand (2014) ³¹	NA	NA	Yes	Unclear	Unclear	Unclear	Unclear	Yes	Unclear	Unclear	Unclear		
Kim (2013) ⁸⁷ (subway drivers)	NA	NA	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Kovacevic (2021) ⁵⁹	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Landolt (2005) ⁹⁴ (PTSS)	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Maeda (2009) ¹⁰⁵ (ship)	Yes	Yes	Yes	Unclear	Unclear	Unclear	Yes	Yes	Yes	Yes	Yes	Unclear	

Table 3. Continued.

	Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	Q9	Q10	Q11	Q12	Q13
Matsuoka (2008) ⁶³	NA	NA	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
Mayou (1997) ⁶⁵	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
McFarlane (1997) ⁶⁶	Yes	Yes	Yes	Unclear	Unclear	Unclear	Yes	Yes	Yes	Yes	Yes	Unclear	
Suliman (2014) ⁷⁸	NA	NA	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	
Mehnert (2012) ⁸⁹	NA	NA	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	
Meiser-Stedman (2009) ⁹⁶	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	
Papadakaki (2017) ⁷²	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	
Naim (2014) ⁶⁷	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	
Nightingale (2000) ⁶⁸	NA	NA	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	
Nishi (2013) ⁷⁰	NA	NA	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	
Mirza (1998) ⁹⁷	Yes	Yes	Yes	Unclear	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	
Pervanidou (2007) ⁹⁸	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	
Ryb (2009) ⁷⁵	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	
Shaikh (2012) ⁷⁷	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	
Smith (2007) ³	Yes	Yes	Yes	Unclear	Unclear	Unclear	Yes	Yes	Yes	Yes	Yes	Unclear	
Stallard (2001) ¹⁰⁰	Yes	Yes	Yes	Unclear	Unclear	Unclear	Yes	Yes	Yes	Yes	Yes	Unclear	
Stallard (2004) ¹⁰¹	Yes	Yes	Yes	Unclear	Unclear	Unclear	Yes	Yes	Yes	Yes	Yes	Unclear	
Sun (2013) ⁷⁹	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	
Lesaca (1996) ⁸⁵	Yes	Yes	Yes	Unclear	Unclear	Unclear	Yes	Yes	Yes	Yes	Yes	Unclear	
Vaiva (2003) ⁸¹	Yes	Yes	Yes	Unclear	Unclear	Unclear	Yes	Yes	Yes	Yes	Yes	Unclear	
Wang (2005) ⁸²	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	
Wu (2016) ¹⁰³	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	
Cross-sectional													
Iteke (2011) ⁵³	Yes	Yes	Yes	Yes	Unclear	Unclear	Yes	Yes					
AlShardan (2020) ²⁵	Yes	Yes	Yes	Yes	Unclear	Unclear	Yes	Yes					
Bedaso (2020) ²⁸	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes					
Berna (2012) ¹⁶	Yes	Yes	Yes	Yes	Unclear	Unclear	Yes	Yes					
Arora (2021) ¹¹⁰	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes					
Khodadadi-Hassankiadeh (2017) ⁵⁷	Yes	Yes	Yes	Yes	Unclear	Unclear	Yes	Yes					
Blanchard (1995) ³⁰	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes					
Bezabh (2018) ²⁹	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes					
Kupchik (2007) ⁶¹	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes					
Ongecha-Owuor (2004) ⁷¹	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes					
Neylan (2021) ¹¹²	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes					
Ning (2017) ⁶⁹	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes					
Yohannes (2018) ⁸⁴	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes					
Williams (2015) ¹⁰²	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes					
Saberi (2013) ⁷⁶	Yes	Yes	Yes	Yes	Unclear	Unclear	Yes	Yes					
Pires (2013) ⁷⁴	Yes	Yes	Yes	Yes	Unclear	Unclear	Yes	Yes					
Prakasam (2013) ⁷³	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes					
Salter (2004) ⁹⁹	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear					
Case-control													
Jones-Alexander (2005) ⁹³	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Unclear		
Ursano (1999) ⁸⁰	Unclear	No	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes		
RCT													
Zehnder (2010) ¹⁰⁴	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Train accident													
Lemos (2018) ⁸⁸	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	
Engelhard (2002) ⁸⁶	Yes	Yes	Yes	Unclear	Unclear	Yes	Yes	Yes	Yes	Yes	Unclear	Yes	

CI: 18.1%-22.8%) among the studied population. We observed a slight variation based on assessment methods, with clinician-administered assessments indicating an 18.7% prevalence (95% CI: 16.0%-21.8%) and self-reported questionnaires showing 22.8% (95% CI: 18.8%-27.3%). After removing outliers, these rates were adjusted to 18.1% (95% CI: 15.4%-21.0%) for clinician-administered assessments and 20.8% (95% CI: 17.5%-24.4%) for self-reported questionnaires. A significant temporal pattern was identified, with PTSD prevalence peaking at 29.4% (95% CI: 22.4%-37.5%) one-month post-RTA, subsequently decreasing to 18.8% (95% CI: 14.8%-23.5%) at three months ($P < 0.001$). Interestingly, age was not a significant predictor of PTSD prevalence rates. We also noted considerable geographic variability in PTSD prevalence, with lower rates observed in Switzerland, Australia, Germany, and Japan, compared to higher rates in Spain, China, and Iran. The included studies were assessed to be of moderate to high quality according to Joanna Briggs Institute standards, ensuring the reliability of these findings. The results of this systematic review and meta-analysis provide a comprehensive overview of the prevalence of PTSD among survivors of RTAs across various geographical regions and assessment methods. The study included data from multiple countries and employed a range of diagnostic tools, resulting in a PTSD prevalence rate ranging from 2.9% to 77.8%. Furthermore, the subgroup analysis revealed that the assessment tools most frequently utilized were PCL-C, CAPS, and SCID. The wide prevalence rate of PTSD might be related to variances in the time interval between the trauma's occurrence and the assessment of PTSD. Factors such as the parameters used to diagnose this disorder and sample characteristics, including gender, type of RTA, and severity of injury, may also influence the PTSD prevalence rates.

Additionally, variations in social support, family stability, and parental involvement may contribute to differences in PTSD prevalence among participants. In a previously published systematic review in 2013, the incidence of PTSD varied from 6 to 45% depending on the type of accident, community support, the severity of the stress, and a history of mental illness.¹⁹ In a recent systematic review of the delayed PTSD prevalence with the current study team, the total prevalence was 13.5%,¹¹³ and in term of risk factors associated with PTSD, Sabahi et al noted that several factors, including female gender, pre-traumatic depression, a history of RTA, peritraumatic dissociative experiences, a diagnosis of ASD, rumination, greater injury severity, and engagement in litigation or compensation following the traumatic event, were significant predictors of PTSD.¹⁸

It is essential to obtain a reliable estimation of PTSD prevalence without adjusting for age following an RTA. This approach will help mental health professionals accurately identify the number of adolescents and children at risk for the disorder and effectively allocate resources

for prevention and treatment interventions. A meta-analysis with a total of 1532 children and adolescents reported that one-fifth of children and adolescents who participated in RTAs later developed PTSD, underscoring the importance of regular PTSD assessments and the implementation of timely psychological interventions for this vulnerable group.¹¹⁴

The adverse psychological repercussions resulting from MVA are significant, with all studies indicating detrimental effects on at least one aspect of psychopathology.¹¹⁵ Failure to address these symptoms may result in a significant risk of progression to severe mental health disorders, including major depressive disorder (MDD), PTSD, panic disorder, and generalized anxiety disorder.^{116,117} A systematic review conducted by Marasini and colleagues revealed a significant prevalence of adverse psychological outcomes after an MVA. The findings demonstrated a consistent pattern, highlighting the predominance of specific psychopathologies, including PTSD, depression, anxiety, travel-related phobia, and emotional distress.¹¹⁵ While Injuries significantly contribute to detrimental mental health outcomes, Individuals who have not entirely recovered from their accidents, along with those who have suffered severe injuries, display a markedly higher likelihood of experiencing adverse psychological effects.^{17,118}

It is imperative to underscore the significance of two prior studies regarding the prevalence and predictors of PTSD among survivors of RTA. The first study conducted by Heron Delaney et al¹⁹ explored various potential predictors of subsequent PTSD following RTA. The authors identified several contributing factors, including rumination related to the traumatic experience, a perceived sense of imminent danger, insufficient social support, the heightened intensity of ASD symptoms, ongoing physical ailments, historical and emotional difficulties, previous anxiety disorders, and participation in legal proceedings or compensation processes, all of which serve as reliable indicators of PTSD.¹⁹ Additionally, a second study conducted by Lin et al presented a meta-analysis encompassing 15 highly heterogeneous studies involving 6,804 RTA survivors, whereby a pooled prevalence rate of 22.25% (95% confidence interval: 16.71%–28.33%) was estimated.¹⁵ Moreover, subgroup analyses indicated that the prevalence of PTSD among RTA survivors exhibited considerable variation across studies, influenced by factors such as the PTSD assessment tool utilized, geographic location, ethnicity, gender, and educational level.

Variability in PTSD Prevalence

Clinician-Administered Measures

The range of PTSD prevalence using these measures is striking, from single-digit percentages to over 80% at some points. While some variation is expected due to differing study populations and methodologies, such extreme differences suggest that other factors are at play. For example, the Lesaca et al⁸⁵ study reported 48% in one

month, and the Kobayashi et al⁵⁸ study reported 81% in six weeks are exceptionally high. These outlier results may be due to unique characteristics of the study samples (e.g., severity of injuries, pre-existing mental health conditions, cultural context) or specific study methodologies. Conversely, studies like Ongecha-Owuor et al,⁷¹ with 13.3% at one month and the study reporting 1.7% at six months, represent the lower end of the spectrum, possibly reflecting more resilient populations or differences in how PTSD was diagnosed.

Self-Report Questionnaires

Similar to clinician-administered measures, self-report data also demonstrates substantial variability. The Allenou et al⁹⁵ study's low prevalence of 3.6% contrasts sharply with the Fekadu et al⁴¹ study's 46.5%. Again, differences in study populations, cultural factors, the specific self-report tool used, and the timing of assessment likely contribute to this variation. The Wang et al⁸² study's extremely high rate of 82.8% at six weeks is a clear outlier that requires further investigation. While the authors suggest a potential link between riding motor scooters and experiencing greater emotional stress (citing the higher prevalence of scooter riders in their sample and Murray et al's¹¹⁹ findings), this alone may not fully account for such a dramatic difference. The Wang et al⁸² study also found a high rate of ASD at one week (72%), which, while similar to Murray et al, was significantly higher than Jaspers' results.¹²⁰ The authors acknowledge that while ASD may be present early on, it doesn't always develop into PTSD. They emphasize the importance of considering various contributing factors, including biological predispositions, pre-existing psychosocial factors, post-accident events, and, crucially, the individual's subjective experience of the trauma. They rightly point out that the subjective meaning of the event could be a powerful predictor of PTSD development. Therefore, while the high prevalence of scooter riders in the Wang et al study⁸² might contribute to their findings, it's likely a complex interplay of factors, including the subjective impact of the accident, that underlies their reported 82.8% PTSD rate.

Our results showed that in the time to measure PTSD between 1 to 6 months after RTAs, the overall prevalence varies across studies, ranging from 8% to 36%. This rate decreased when we performed a subgroup analysis on the studies that evaluated PTSD between 4 and 6 months after RTA, varying from 18% to 28%. However, we should consider that only limited studies were eligible to be included in the latter subgroup analysis compared to those in the first subgroup. In addition, to determine the prevalence of PTSD, all related studies on whether cases were admitted to the emergency department and outpatients were included in the meta-analysis. The analysis results showed that a higher prevalence of PTSD was observed in cases with serious injury (20.3%). In comparison, 31% of hospitalized patients following RTA met PTSD criteria at the time point of 1-6 months.

The rate was 20% for those attending the emergency department cases.

The present study systematically assessed the prevalence among adult and adolescent survivors of RTAs. The results indicated older RTA victims (21.5%) were more susceptible to PTSD than younger ones (17.4%). Evidence suggests that age-related stressors can intensify PTSD symptoms. Factors such as role and functional changes accompanying aging—retirement, bereavement, declining physical health that limits autonomy, and reduced social support may contribute to an increase in PTSD symptoms.¹²¹ A comprehensive subgroup analysis was conducted, considering various temporal dimensions across distinct age categories and the severity of injuries—differentiating between severe and life-threatening injuries versus minor injuries and outpatient conditions.

Here, we noticed the difference between self-reported (20.3%) and clinician-based PTSD prevalence (22.8%) results that stem from various factors. Self-reports can be influenced by subjective biases, emotional states, or limited understanding of PTSD symptoms, leading to over- or underestimation. Clinicians, by contrast, use structured interviews and standardized diagnostic criteria, providing greater precision. Survivors may interpret symptoms differently in self-reports or struggle to distinguish PTSD from other conditions. Additionally, rapport with clinicians can affect disclosure during interviews, while cultural or language barriers may influence the accuracy of both methods. Severe symptoms can also impair insight, affecting self-reports, whereas clinicians can contextualize symptoms more effectively. Combining both approaches enhances diagnostic accuracy and understanding.¹²²⁻¹²⁴

Additionally, the analysis was stratified by World Health Organization (WHO) regions and specific countries to enhance the contextual understanding of PTSD prevalence. However, this study faced some limitations. Notably, it did not incorporate research with delayed PTSD assessments. This might have implications for the comprehensiveness of our findings. Furthermore, the lack of a gender-specific subgroup analysis limits the capacity to make detailed conclusions about how gender affects PTSD outcomes. Furthermore, because few studies report PTSD rates among survivors based on their position in the vehicle, analyzing this variable was not possible.

Recommendations for Future Studies

Based on the findings of this study, several clinical recommendations can be made to enhance the identification and management of PTSD among RTA survivors. Firstly, it is crucial to implement routine screening for PTSD among RTA survivors. The early utilization of validated assessment tools, particularly during the initial months following the accident, facilitates the timely identification of high-risk individuals. However, the precise implication of therapeutic interventions in self-reported PTSD victims is necessary to avoid overdiagnosis and overtreatment. Furthermore, practical psychological

support services should be integrated into treatment plans, encompassing Cognitive Behavioral Therapy and family counseling. Given the variability of PTSD prevalence rates and demographic factors, interventions must be tailored accordingly. Healthcare providers need to be adequately developed to enhance their understanding of PTSD and its potential manifestations in survivors, particularly at the primary level of care.

To enhance the understanding of PTSD among RTA survivors, future studies should concentrate on various demographic groups. Given the limited research on PTSD in children and adolescents, they should prioritize these populations. Then, more epidemiological research can clarify the explicit causal relationships between factors, leading to targeted preventive strategies. By examining the relationship between PTSD and other psychological disorders like depression and anxiety, valuable insights will be gained for developing treatment strategies that encompass various aspects of mental health in survivors of RTAs. Studies should evaluate the effectiveness and differences between self-reported and clinically administered approaches to suggest a reliable and optimized strategy. Finally, the enhanced understanding and management of survivors lead to improved recovery and quality of life.

Conclusion

The prevalence of PTSD among survivors was measured at 20.3%. Countries such as Switzerland, Australia, Germany, and Japan exhibited the lowest prevalence rates, whereas Spain, China, and Iran recorded the highest prevalence rates. Moreover, the subgroup analysis indicated that the frequently employed checklists and criteria for the assessment of PTSD included the PCL-C, the CAPS, and the SCID. The severity of injuries sustained significantly influenced the prevalence of PTSD. Future epidemiological studies are warranted to investigate potential causal relationships between the positioning of individuals within vehicles and the development of PTSD. This exploration aims to enhance our comprehension of the determinants of PTSD and subsequently improve prevention and treatment strategies for survivors of RTAs.

Acknowledgments

The authors would like to thank the regional ethics committee of Tabriz University of Medical Sciences for approving the proposal.

Authors' Contribution

Conceptualization: Homayoun Sadeghi-Bazargani, Kavous Shahsavarinia, Sakineh Hajebrabimi.

Data curation: Zahra Sabahi, Haleh Farshi, Fateme Tahmasbi, Nooshin Milanchian, Mostafa Farahbakhsh.

Formal analysis: Hanieh Salehi-Pourmehr.

Funding acquisition: Hanieh Salehi-Pourmehr.

Investigation: Zahra Sabahi, Homayoun Sadeghi-Bazargani, Kavous Shahsavarinia.

Methodology: Hanieh Salehi-Pourmehr, Sakineh Hajebrabimi, Javad Mahmoudi.

Project administration: Zahra Sabahi, Haleh Farshi, Fateme Tahmasbi, Nooshin Milanchian.

Resources: Haleh Farshi, Fateme Tahmasbi, Nooshin Milanchian.

Software: Hanieh Salehi-Pourmehr.

Supervision: Hanieh Salehi-Pourmehr, Homayoun Sadeghi-Bazargani.

Validation: Homayoun Sadeghi-Bazargani, Kavous Shahsavarinia.

Visualization: Hanieh Salehi-Pourmehr, Homayoun Sadeghi-Bazargani, Kavous Shahsavarinia, Javad Mahmoudi.

Writing-original draft: Zahra Sabahi, Haleh Farshi, Fateme Tahmasbi, Nooshin Milanchian, Mostafa Farahbakhsh, Javad Mahmoudi.

Writing-review & editing: Homayoun Sadeghi-Bazargani, Kavous Shahsavarinia, Sakineh Hajebrabimi, Hanieh Salehi-Pourmehr.

Competing Interests

None.

Data Availability Statement

Not applicable.

Ethical Approval

The regional ethics committee of Tabriz University of Medical Sciences approved this study (IR.TBZMED.REC.1400.966).

Funding

This study is supported by the Road Traffic Injury Research Center, Tabriz University of Medical Sciences, Tabriz, Iran (Grant No: 68826).

Supplementary Files

Supplementary file 1. Search strategy of PubMed (PTSD+Traffic Accident).

Supplementary file 2 contains Figures S1-S11.

References

1. World Health Organization (WHO). Global Status Report on Road Safety 2018. Available from: <https://www.who.int/publications/i/item/9789241565684>.
2. Asuquo JE, Edet BE, Abang IE, Essien EA, Osakwe OG, Aigbomain EJ, et al. Depression and posttraumatic stress disorder among road traffic accident victims managed in a tertiary hospital in Southern Nigeria. *Niger J Clin Pract*. 2017;20(2):170-5. doi: 10.4103/1119-3077.196114.
3. Smith B, Mackenzie-Ross S, Scragg P. Prevalence of poor psychological morbidity following a minor road traffic accident (RTA): the clinical implications of a prospective longitudinal study. *Couns Psychol Q*. 2007;20(2):149-55. doi: 10.1080/09515070701403679.
4. Hamanaka S, Asukai N, Kamijo Y, Hatta K, Kishimoto J, Miyaoka H. Acute stress disorder and posttraumatic stress disorder symptoms among patients severely injured in motor vehicle accidents in Japan. *Gen Hosp Psychiatry*. 2006;28(3):234-41. doi: 10.1016/j.genhosppsych.2006.02.007.
5. Blanchard EB, Hickling EJ. After the Crash: Psychological Assessment and Treatment of Survivors of Motor Vehicle Accidents. American Psychological Association; 2004. doi: 10.1037/10676-000.
6. National Center for Statistics and Analytics. Traffic Safety Quick Facts 2017 (DOT HS 812 747). Washington, DC: National Highway Traffic Safety Administration.
7. Schäfer I, Barkmann C, Riedesser P, Schulte-Markwort M. Posttraumatic syndromes in children and adolescents after road traffic accidents--a prospective cohort study. *Psychopathology*. 2006;39(4):159-64. doi: 10.1159/000092676.
8. Tierens M, Bal S, Crombez G, Van de Voorde P, Rosseel Y, Antrop I, et al. The traumatic impact of motor vehicle accidents in high school students. *J Pediatr Psychol*. 2012;37(1):1-10. doi: 10.1093/jpepsy/jsr058.
9. Stallard P, Velleman R, Baldwin S. Prospective study of post-traumatic stress disorder in children involved in road traffic

- accidents. *BMJ*. 1998;317(7173):1619-23. doi: [10.1136/bmj.317.7173.1619](#).
10. American Psychological Association (APA). *Diagnostic and Statistical Manual of Mental Disorders*. 5th ed. APA; 2013. doi: [10.1176/appi.books.9780890425596](#).
 11. Spitzer R, Williams J. *Diagnostic and Statistical Manual of Mental Disorders*. revised. DSM III R. Washington, DC: American Psychiatric Association (APA); 1987. p. 47-217.
 12. Dai W, Kaminga AC, Tan H, Wang J, Lai Z, Wu X, et al. Comorbidity of post-traumatic stress disorder and anxiety in flood survivors: prevalence and shared risk factors. *Medicine (Baltimore)*. 2017;96(36):e7994. doi: [10.1097/md.00000000000007994](#).
 13. Landolt MA, Vollrath ME, Gnehm HE, Sennhauser FH. Post-traumatic stress impacts on quality of life in children after road traffic accidents: prospective study. *Aust N Z J Psychiatry*. 2009;43(8):746-53. doi: [10.1080/00048670903001919](#).
 14. Bartoli F, Crocamo C, Alamia A, Amidani F, Paggi E, Pini E, et al. Posttraumatic stress disorder and risk of obesity: systematic review and meta-analysis. *J Clin Psychiatry*. 2015;76(10):e1253-61. doi: [10.4088/JCP.14r09199](#).
 15. Lin W, Gong L, Xia M, Dai W. Prevalence of posttraumatic stress disorder among road traffic accident survivors: a PRISMA-compliant meta-analysis. *Medicine (Baltimore)*. 2018;97(3):e9693. doi: [10.1097/md.00000000000009693](#).
 16. Berna G, Vaiva G, Ducrocq F, Duhem S, Nandrino JL. Categorical and dimensional study of the predictive factors of the development of a psychotrauma in victims of car accidents. *J Anxiety Disord*. 2012;26(1):239-45. doi: [10.1016/j.janxdis.2011.11.011](#).
 17. Pozzato I, Kifley A, Craig A, Gopinath B, Tran Y, Jagnoor J, et al. Effects of seeking compensation on the psychological health and recovery of injured patients: the role of stress vulnerability and injury-related disability. *Psychol Med*. 2022;52(1):68-79. doi: [10.1017/s003329172000166x](#).
 18. Sabahi Z, Hasani P, Salehi-Pourmehr H, Beheshti R, Sadeghi-Bazargani H. What are the predictors of post-traumatic stress disorder among road traffic accident survivors: a systematic review. *J Nerv Ment Dis*. 2024;212(2):104-16. doi: [10.1097/nmd.0000000000001739](#).
 19. Heron-Delaney M, Kenardy J, Charlton E, Matsuoka Y. A systematic review of predictors of posttraumatic stress disorder (PTSD) for adult road traffic crash survivors. *Injury*. 2013;44(11):1413-22. doi: [10.1016/j.injury.2013.07.011](#).
 20. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71. doi: [10.1136/bmj.n71](#).
 21. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med*. 2002;21(11):1539-58. doi: [10.1002/sim.1186](#).
 22. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997;315(7109):629-34. doi: [10.1136/bmj.315.7109.629](#).
 23. Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics*. 1994;50(4):1088-101.
 24. Singh JA, Siddiqi M, Parameshwar P, Chandra-Mouli V. World Health Organization guidance on ethical considerations in planning and reviewing research studies on sexual and reproductive health in adolescents. *J Adolesc Health*. 2019;64(4):427-9. doi: [10.1016/j.jadohealth.2019.01.008](#).
 25. Alshardan AM, Alshuqaybi HM, Agha S, Masuadi E, Albabtain I, Albugami AS, et al. Assessment of sleep quality, post-traumatic stress disorder, and locus of control in motor vehicle accident survivors. *Australas Med J*. 2020;13(3):78-85. doi: [10.35841/1836-1935.13.3.78-85](#).
 26. Angerpointner K, Weber S, Tschach K, Schubert H, Herbst T, Ernstberger A, et al. Posttraumatic stress disorder after minor trauma - a prospective cohort study. *Med Hypotheses*. 2020;135:109465. doi: [10.1016/j.mehy.2019.109465](#).
 27. Bahari R, Mohamad Alwi MN, Ahmad MR, Mohd Saiboon I. Incidence and demographical characteristics of patients with post-traumatic stress disorder due to motor vehicle accidents. *Malays Fam Physician*. 2017;12(3):2-7.
 28. Bedaso A, Kediro G, Ebrahim J, Tadesse F, Mekonnen S, Gobena N, et al. Prevalence and determinants of post-traumatic stress disorder among road traffic accident survivors: a prospective survey at selected hospitals in southern Ethiopia. *BMC Emerg Med*. 2020;20(1):52. doi: [10.1186/s12873-020-00348-5](#).
 29. Bezabh YH, Abebe SM, Fanta T, Tadesse A, Tulu M. Prevalence and associated factors of post-traumatic stress disorder among emergency responders of Addis Ababa Fire and Emergency Control and Prevention Service Authority, Ethiopia: institution-based, cross-sectional study. *BMJ Open*. 2018;8(7):e020705. doi: [10.1136/bmjopen-2017-020705](#).
 30. Blanchard EB, Hickling EJ, Taylor AE, Loos W. Psychiatric morbidity associated with motor vehicle accidents. *J Nerv Ment Dis*. 1995;183(8):495-504. doi: [10.1097/00005053-199508000-00001](#).
 31. Brand S, Otte D, Petri M, Decker S, Stübiger T, Krettek C, et al. Incidence of posttraumatic stress disorder after traffic accidents in Germany. *Int J Emerg Ment Health*. 2014;16(1):233-6. doi: [10.4172/1522-4821.1000109](#).
 32. Bryant RA, Harvey AG, Guthrie RM, Moulds ML. Acute psychophysiological arousal and posttraumatic stress disorder: a two-year prospective study. *J Trauma Stress*. 2003;16(5):439-43. doi: [10.1023/a:1025750209553](#).
 33. Bryant RA, Harvey AG, Guthrie RM, Moulds ML. A prospective study of psychophysiological arousal, acute stress disorder, and posttraumatic stress disorder. *J Abnorm Psychol*. 2000;109(2):341-4.
 34. Buckley B, Nugent N, Sledjeski E, Raimonde AJ, Spoonster E, Bogart LM, et al. Evaluation of initial posttrauma cardiovascular levels in association with acute PTSD symptoms following a serious motor vehicle accident. *J Trauma Stress*. 2004;17(4):317-24. doi: [10.1023/B:JOTS.0000038480.87290.4a](#).
 35. Chossegros L, Hours M, Charnay P, Bernard M, Fort E, Boisson D, et al. Predictive factors of chronic post-traumatic stress disorder 6 months after a road traffic accident. *Accid Anal Prev*. 2011;43(1):471-7. doi: [10.1016/j.aap.2010.10.004](#).
 36. Coronas R, Gallardo O, Moreno MJ, Suárez D, García-Parés G, Menchón JM. Heart rate measured in the acute aftermath of trauma can predict post-traumatic stress disorder: a prospective study in motor vehicle accident survivors. *Eur Psychiatry*. 2011;26(8):508-12. doi: [10.1016/j.eurpsy.2010.06.006](#).
 37. Delahanty DL, Royer DK, Raimonde AJ, Spoonster E. Peritraumatic dissociation is inversely related to catecholamine levels in initial urine samples of motor vehicle accident victims. *J Trauma Dissociation*. 2003;4(1):65-80. doi: [10.1300/J229v04n01_05](#).
 38. Doohan I, Björnstig U, Östlund U, Saveman BI. Exploring injury panorama, consequences, and recovery among bus crash survivors: a mixed-methods research study. *Prehosp Disaster Med*. 2017;32(2):165-74. doi: [10.1017/s1049023x16001485](#).
 39. Ehlers A, Mayou RA, Bryant B. Psychological predictors of chronic posttraumatic stress disorder after motor vehicle accidents. *J Abnorm Psychol*. 1998;107(3):508-19. doi: [10.1037//0021-843x.107.3.508](#).
 40. Ehling T, Ehlers A, Glucksman E. Do cognitive models help in predicting the severity of posttraumatic stress disorder, phobia, and depression after motor vehicle accidents? A prospective longitudinal study. *J Consult Clin Psychol*. 2008;76(2):219-30. doi: [10.1037/0022-006x.76.2.219](#).
 41. Fekadu W, Mekonen T, Belete H, Belete A, Yohannes K.

- Incidence of post-traumatic stress disorder after road traffic accident. *Front Psychiatry*. 2019;10:519. doi: [10.3389/fpsyt.2019.00519](https://doi.org/10.3389/fpsyt.2019.00519).
42. Fitzharris M, Fildes B, Charlton J. Anxiety, acute- and post-traumatic stress symptoms following involvement in traffic crashes. *Annu Proc Assoc Adv Automot Med*. 2006;50:297-315.
 43. Flesher MR, Delahanty DL, Raimonde AJ, Spoonster E. Amnesia, neuroendocrine levels and PTSD in motor vehicle accident victims. *Brain Inj*. 2001;15(10):879-89. doi: [10.1080/02699050110065682](https://doi.org/10.1080/02699050110065682).
 44. Fredman SJ, Beck JG, Shnaider P, Le Y, Pukay-Martin ND, Pentel KZ, et al. Longitudinal associations between PTSD symptoms and dyadic conflict communication following a severe motor vehicle accident. *Behav Ther*. 2017;48(2):235-46. doi: [10.1016/j.beth.2016.05.001](https://doi.org/10.1016/j.beth.2016.05.001).
 45. Frommberger UH, Stieglitz RD, Nyberg E, Schlickewei W, Kuner E, Berger M. Prediction of posttraumatic stress disorder by immediate reactions to trauma: a prospective study in road traffic accident victims. *Eur Arch Psychiatry Clin Neurosci*. 1998;248(6):316-21. doi: [10.1007/s004060050057](https://doi.org/10.1007/s004060050057).
 46. Fuglsang AK, Moergeli H, Schnyder U. Does acute stress disorder predict post-traumatic stress disorder in traffic accident victims? Analysis of a self-report inventory. *Nord J Psychiatry*. 2004;58(3):223-9. doi: [10.1080/08039480410006304](https://doi.org/10.1080/08039480410006304).
 47. Fullerton CS, Ursano RJ, Epstein RS, Crowley B, Vance K, Kao TC, et al. Gender differences in posttraumatic stress disorder after motor vehicle accidents. *Am J Psychiatry*. 2001;158(9):1486-91. doi: [10.1176/appi.ajp.158.9.1486](https://doi.org/10.1176/appi.ajp.158.9.1486).
 48. Gabert-Quillen CA, Irish LA, Sledjeski E, Fallon W, Spoonster E, Delahanty DL. The impact of social support on the relationship between trauma history and PTSD symptoms in motor vehicle accident victims. *Int J Stress Manag*. 2012;19(1):69-79. doi: [10.1037/a0026488](https://doi.org/10.1037/a0026488).
 49. Harvey AG, Bryant RA. The relationship between acute stress disorder and posttraumatic stress disorder: a prospective evaluation of motor vehicle accident survivors. *J Consult Clin Psychol*. 1998;66(3):507-12. doi: [10.1037//0022-006x.66.3.507](https://doi.org/10.1037//0022-006x.66.3.507).
 50. Holeva V, Tarrier N, Wells A. Prevalence and predictors of acute stress disorder and PTSD following road traffic accidents: thought control strategies and social support. *Behav Ther*. 2001;32(1):65-83. doi: [10.1016/s0005-7894\(01\)80044-7](https://doi.org/10.1016/s0005-7894(01)80044-7).
 51. Hu H, Sun Y, Su S, Wang Y, Qiu Y, Yang X, et al. Cortical surface area reduction in identification of subjects at high risk for post-traumatic stress disorder: a pilot study. *Aust N Z J Psychiatry*. 2018;52(11):1084-91. doi: [10.1177/0004867417750757](https://doi.org/10.1177/0004867417750757).
 52. Irish LA, Fischer B, Fallon W, Spoonster E, Sledjeski EM, Delahanty DL. Gender differences in PTSD symptoms: an exploration of peritraumatic mechanisms. *J Anxiety Disord*. 2011;25(2):209-16. doi: [10.1016/j.janxdis.2010.09.004](https://doi.org/10.1016/j.janxdis.2010.09.004).
 53. Iteke O, Bakare MO, Agomoh AO, Uwakwe R, Onwukwe JU. Road traffic accidents and posttraumatic stress disorder in an orthopedic setting in south-eastern Nigeria: a controlled study. *Scand J Trauma Resusc Emerg Med*. 2011;19:39. doi: [10.1186/1757-7241-19-39](https://doi.org/10.1186/1757-7241-19-39).
 54. Jeavons S. Predicting who suffers psychological trauma in the first year after a road accident. *Behav Res Ther*. 2000;38(5):499-508. doi: [10.1016/s0005-7967\(99\)00073-x](https://doi.org/10.1016/s0005-7967(99)00073-x).
 55. Jones C, Harvey AG, Brewin CR. Traumatic brain injury, dissociation, and posttraumatic stress disorder in road traffic accident survivors. *J Trauma Stress*. 2005;18(3):181-91. doi: [10.1002/jts.20031](https://doi.org/10.1002/jts.20031).
 56. Kassam-Adams N, Fleisher CL, Winston FK. Acute stress disorder and posttraumatic stress disorder in parents of injured children. *J Trauma Stress*. 2009;22(4):294-302. doi: [10.1002/jts.20424](https://doi.org/10.1002/jts.20424).
 57. Khodadadi-Hassankiadeh N, Dehghan Nayeri N, Shahsavari H, Yousefzadeh-Chabok S, Haghani H. Predictors of post-traumatic stress disorder among victims of serious motor vehicle accidents. *Int J Community Based Nurs Midwifery*. 2017;5(4):355-64.
 58. Kobayashi I, Sledjeski EM, Delahanty DL. Gender and age interact to predict the development of posttraumatic stress disorder symptoms following a motor vehicle accident. *Psychol Trauma*. 2019;11(3):328-36. doi: [10.1037/tra0000366](https://doi.org/10.1037/tra0000366).
 59. Kovacevic J, Fotez I, Miskulin I, Lesic D, Miskulin M, Berlancic T, et al. Different patterns of mental health outcomes among road traffic crash survivors: a prospective cohort study. *Int J Environ Res Public Health*. 2021;18(4):1564. doi: [10.3390/ijerph18041564](https://doi.org/10.3390/ijerph18041564).
 60. Kühn M, Ehler U, Rumpf HJ, Backhaus J, Hohagen F, Broocks A. Onset and maintenance of psychiatric disorders after serious accidents. *Eur Arch Psychiatry Clin Neurosci*. 2006;256(8):497-503. doi: [10.1007/s00406-006-0670-6](https://doi.org/10.1007/s00406-006-0670-6).
 61. Kupchik M, Strous RD, Erez R, Gonen N, Weizman A, Spivak B. Demographic and clinical characteristics of motor vehicle accident victims in the community general health outpatient clinic: a comparison of PTSD and non-PTSD subjects. *Depress Anxiety*. 2007;24(4):244-50. doi: [10.1002/da.20189](https://doi.org/10.1002/da.20189).
 62. Li X, Sun L, Li Q, Wang L. Prediction of posttraumatic stress disorder by acute stress disorder in traffic accident survivors. *Turk J Med Sci*. 2021;51(5):2502-9. doi: [10.3906/sag-2008-282](https://doi.org/10.3906/sag-2008-282).
 63. Matsuoka Y, Nishi D, Nakajima S, Kim Y, Homma M, Otomo Y. Incidence and prediction of psychiatric morbidity after a motor vehicle accident in Japan: the Tachikawa Cohort of Motor Vehicle Accident Study. *Crit Care Med*. 2008;36(1):74-80. doi: [10.1097/01.Ccm.0000291650.70816.D6](https://doi.org/10.1097/01.Ccm.0000291650.70816.D6).
 64. Mayou R, Bryant B, Duthie R. Psychiatric consequences of road traffic accidents. *BMJ*. 1993;307(6905):647-51. doi: [10.1136/bmj.307.6905.647](https://doi.org/10.1136/bmj.307.6905.647).
 65. Mayou R, Tyndel S, Bryant B. Long-term outcome of motor vehicle accident injury. *Psychosom Med*. 1997;59(6):578-84. doi: [10.1097/00006842-199711000-00004](https://doi.org/10.1097/00006842-199711000-00004).
 66. McFarlane AC, Atchison M, Yehuda R. The acute stress response following motor vehicle accidents and its relation to PTSD. *Ann N Y Acad Sci*. 1997;821:437-41. doi: [10.1111/j.1749-6632.1997.tb48299.x](https://doi.org/10.1111/j.1749-6632.1997.tb48299.x).
 67. Naim R, Wald I, Lior A, Pine DS, Fox NA, Sheppes G, et al. Perturbed threat monitoring following a traumatic event predicts risk for post-traumatic stress disorder. *Psychol Med*. 2014;44(10):2077-84. doi: [10.1017/s0033291713002456](https://doi.org/10.1017/s0033291713002456).
 68. Nightingale J, Williams RM. Attitudes to emotional expression and personality in predicting post-traumatic stress disorder. *Br J Clin Psychol*. 2000;39(3):243-54. doi: [10.1348/014466500163266](https://doi.org/10.1348/014466500163266).
 69. Ning L, Guan S, Liu J. Impact of personality and social support on posttraumatic stress disorder after traffic accidents. *Medicine (Baltimore)*. 2017;96(34):e7815. doi: [10.1097/md.0000000000007815](https://doi.org/10.1097/md.0000000000007815).
 70. Nishi D, Noguchi H, Yonemoto N, Nakajima S, Kim Y, Matsuoka Y. Incidence and prediction of post-traumatic stress disorder at 6 months after motor vehicle accidents in Japan. *Psychosomatics*. 2013;54(3):263-71. doi: [10.1016/j.psych.2012.07.010](https://doi.org/10.1016/j.psych.2012.07.010).
 71. Ongecha-Owuor FA, Kathuku DM, Othieno CJ, Ndeti DM. Post-traumatic stress disorder among motor vehicle accident survivors attending the orthopaedic and trauma clinic at Kenyatta National Hospital, Nairobi. *East Afr Med J*. 2004;81(7):362-6. doi: [10.4314/eamj.v81i7.9192](https://doi.org/10.4314/eamj.v81i7.9192).
 72. Papadakaki M, Ferraro OE, Orsi C, Otte D, Tzamalouka G, von-der-Geest M, et al. Psychological distress and physical disability in patients sustaining severe injuries in road traffic crashes: results from a one-year cohort study from three European countries. *Injury*. 2017;48(2):297-306. doi: [10.1016/j.injury.2016.11.011](https://doi.org/10.1016/j.injury.2016.11.011).
 73. Prakasam A, Joseph A, Zachariah N, Joy S. Assess the impact of

- post-traumatic stress disorder in road traffic accident survivors in Erode district, Tamil Nadu. *Int J Pharm Res.* 2020;12(1):109-15. doi: [10.31838/ijpr/2020.12.01.009](#).
74. Pires TS, da Costa Maia A. Posttraumatic stress disorder among serious motor vehicle accident victims: analysis of predictors. *Rev Psiquiatr Clin.* 2013;40(6):211-4.
 75. Ryb GE, Dischinger PC, Read KM, Kufera JA. PTSD after severe vehicular crashes. *Ann Adv Automot Med.* 2009;53:177-93.
 76. Saberi HR, Abbasian H, Motalebi Kashani M, Naseri Esfahani AH. Post-traumatic stress disorder: a neglected health concern among commercial motor vehicle drivers. *Int J Occup Environ Med.* 2013;4(4):185-94.
 77. Shaikh Al Arab A, Guédon-Moreau L, Ducrocq F, Molenda S, Duhem S, Salleron J, et al. Temporal analysis of heart rate variability as a predictor of post-traumatic stress disorder in road traffic accidents survivors. *J Psychiatr Res.* 2012;46(6):790-6. doi: [10.1016/j.jpsychires.2012.02.006](#).
 78. Suliman S, Stein DJ, Seedat S. Clinical and neuropsychological predictors of posttraumatic stress disorder. *Medicine (Baltimore).* 2014;93(22):e113. doi: [10.1097/md.0000000000000113](#).
 79. Sun Y, Wang Z, Ding W, Wan J, Zhuang Z, Zhang Y, et al. Alterations in white matter microstructure as vulnerability factors and acquired signs of traffic accident-induced PTSD. *PLoS One.* 2013;8(12):e83473. doi: [10.1371/journal.pone.0083473](#).
 80. Ursano RJ, Fullerton CS, Epstein RS, Crowley B, Kao TC, Vance K, et al. Acute and chronic posttraumatic stress disorder in motor vehicle accident victims. *Am J Psychiatry.* 1999;156(4):589-95. doi: [10.1176/ajp.156.4.589](#).
 81. Vaiva G, Brunet A, Lebigot F, Boss V, Ducrocq F, Devos P, et al. Fright (effroi) and other peritraumatic responses after a serious motor vehicle accident: prospective influence on acute PTSD development. *Can J Psychiatry.* 2003;48(6):395-401. doi: [10.1177/070674370304800606](#).
 82. Wang CH, Tsay SL, Bond AE. Post-traumatic stress disorder, depression, anxiety and quality of life in patients with traffic-related injuries. *J Adv Nurs.* 2005;52(1):22-30. doi: [10.1111/j.1365-2648.2005.03560.x](#).
 83. Yaşan A, Guzel A, Tamam Y, Ozkan M. Predictive factors for acute stress disorder and posttraumatic stress disorder after motor vehicle accidents. *Psychopathology.* 2009;42(4):236-41. doi: [10.1159/000218521](#).
 84. Yohannes K, Gebeyehu A, Adera T, Ayano G, Fekadu W. Prevalence and correlates of post-traumatic stress disorder among survivors of road traffic accidents in Ethiopia. *Int J Ment Health Syst.* 2018;12:50. doi: [10.1186/s13033-018-0229-8](#).
 85. Lesaca T. Symptoms of stress disorder and depression among trauma counselors after an airline disaster. *Psychiatr Serv.* 1996;47(4):424-6. doi: [10.1176/ps.47.4.424](#).
 86. Engelhard IM, van den Hout MA, Arntz A, McNally RJ. A longitudinal study of "intrusion-based reasoning" and posttraumatic stress disorder after exposure to a train disaster. *Behav Res Ther.* 2002;40(12):1415-24. doi: [10.1016/s0005-7967\(02\)00018-9](#).
 87. Kim HR, Yim HW, Jo SJ, Choi B, Jeong SH, Lee KS, et al. Major depressive disorder, panic disorder, and post-traumatic stress disorder in Korean subway drivers. *Int Arch Occup Environ Health.* 2013;86(4):471-7. doi: [10.1007/s00420-012-0779-z](#).
 88. da Conceição Lemos I, Patrão I. Train accidents: development of post-traumatic stress disorder in train drivers. *Psychol Commun Health.* 2018;7(1):44-56. doi: [10.5964/pch.v7i1.184](#).
 89. Mehnert A, Nanninga I, Fauth M, Schäfer I. Course and predictors of posttraumatic stress among male train drivers after the experience of 'person under the train' incidents. *J Psychosom Res.* 2012;73(3):191-6. doi: [10.1016/j.jpsychores.2012.06.007](#).
 90. Bryant B, Mayou R, Wiggs L, Ehlers A, Stores G. Psychological consequences of road traffic accidents for children and their mothers. *Psychol Med.* 2004;34(2):335-46. doi: [10.1017/s0033291703001053](#).
 91. Di Gallo A, Barton J, Parry-Jones WL. Road traffic accidents: early psychological consequences in children and adolescents. *Br J Psychiatry.* 1997;170:358-62. doi: [10.1192/bjp.170.4.358](#).
 92. Gillies ML, Barton J, Di Gallo A. Follow-up of young road accident victims. *J Trauma Stress.* 2003;16(5):523-6. doi: [10.1023/a:1025774915005](#).
 93. Jones-Alexander J, Blanchard EB, Hickling EJ. Psychophysiological assessment of youthful motor vehicle accident survivors. *Appl Psychophysiol Biofeedback.* 2005;30(2):115-23. doi: [10.1007/s10484-005-4307-8](#).
 94. Landolt MA, Vollrath M, Timm K, Gnehm HE, Sennhauser FH. Predicting posttraumatic stress symptoms in children after road traffic accidents. *J Am Acad Child Adolesc Psychiatry.* 2005;44(12):1276-83. doi: [10.1097/01.chi.0000181045.13960.67](#).
 95. Allenou C, Oliac B, Bourdet-Loubère S, Brunet A, David AC, Claudet I, et al. Symptoms of traumatic stress in mothers of children victims of a motor vehicle accident. *Depress Anxiety.* 2010;27(7):652-7. doi: [10.1002/da.20650](#).
 96. Meiser-Stedman R, Dalgleish T, Glucksman E, Yule W, Smith P. Maladaptive cognitive appraisals mediate the evolution of posttraumatic stress reactions: a 6-month follow-up of child and adolescent assault and motor vehicle accident survivors. *J Abnorm Psychol.* 2009;118(4):778-87. doi: [10.1037/a0016945](#).
 97. Mirza KA, Bhadrinath BR, Goodyer IM, Gilmour C. Post-traumatic stress disorder in children and adolescents following road traffic accidents. *Br J Psychiatry.* 1998;172:443-7. doi: [10.1192/bjp.172.5.443](#).
 98. Pervanidou P, Kolaitis G, Charitaki S, Margeli A, Ferentinos S, Bakoula C, et al. Elevated morning serum interleukin (IL)-6 or evening salivary cortisol concentrations predict posttraumatic stress disorder in children and adolescents six months after a motor vehicle accident. *Psychoneuroendocrinology.* 2007;32(8-10):991-9. doi: [10.1016/j.psyneuen.2007.07.001](#).
 99. Salter E, Stallard P. Posttraumatic growth in child survivors of a road traffic accident. *J Trauma Stress.* 2004;17(4):335-40. doi: [10.1023/b:jots.0000038482.53911.01](#).
 100. Stallard P, Velleman R, Langsford J, Baldwin S. Coping and psychological distress in children involved in road traffic accidents. *Br J Clin Psychol.* 2001;40(2):197-208. doi: [10.1348/014466501163643](#).
 101. Stallard P, Salter E, Velleman R. Posttraumatic stress disorder following road traffic accidents - a second prospective study. *Eur Child Adolesc Psychiatry.* 2004;13(3):172-8. doi: [10.1007/s00787-004-0399-6](#).
 102. Williams JL, Rheingold AA, Knowlton AW, Saunders BE, Kilpatrick DG. Associations between motor vehicle crashes and mental health problems: data from the National Survey of Adolescents-Replication. *J Trauma Stress.* 2015;28(1):41-8. doi: [10.1002/jts.21983](#).
 103. Wu F, Meng WY, Hao CZ, Zhu LL, Chen DQ, Lin LY, et al. Analysis of posttraumatic stress disorder in children with road traffic injury in Wenzhou, China. *Traffic Inj Prev.* 2016;17(2):159-63. doi: [10.1080/15389588.2015.1050719](#).
 104. Zehnder D, Meuli M, Landolt MA. Effectiveness of a single-session early psychological intervention for children after road traffic accidents: a randomised controlled trial. *Child Adolesc Psychiatry Ment Health.* 2010;4:7. doi: [10.1186/1753-2000-4-7](#).
 105. Maeda M, Kato H, Maruoka T. Adolescent vulnerability to PTSD and effects of community-based intervention: longitudinal study among adolescent survivors of the Ehime Maru sea accident. *Psychiatry Clin Neurosci.* 2009;63(6):747-53. doi: [10.1111/j.1440-1819.2009.02031.x](#).

106. Giannopoulou I, Richardson C, Papadatou D. Peer loss: posttraumatic stress, depression, and grief symptoms in a traumatized adolescent community. *Clin Child Psychol Psychiatry*. 2021;26(2):556-68. doi: [10.1177/1359104520980028](https://doi.org/10.1177/1359104520980028).
107. Ziobrowski HN, Kennedy CJ, Ustun B, House SL, Beaudoin FL, An X, et al. Development and validation of a model to predict posttraumatic stress disorder and major depression after a motor vehicle collision. *JAMA Psychiatry*. 2021;78(11):1228-37. doi: [10.1001/jamapsychiatry.2021.2427](https://doi.org/10.1001/jamapsychiatry.2021.2427).
108. Kessler RC, Ressler KJ, House SL, Beaudoin FL, An X, Stevens JS, et al. Socio-demographic and trauma-related predictors of PTSD within 8 weeks of a motor vehicle collision in the AURORA study. *Mol Psychiatry*. 2021;26(7):3108-21. doi: [10.1038/s41380-020-00911-3](https://doi.org/10.1038/s41380-020-00911-3).
109. Joormann J, Ziobrowski HN, King AJ, Gildea SM, Lee S, Sampson NA, et al. Prior histories of posttraumatic stress disorder and major depression and their onset and course in the three months after a motor vehicle collision in the AURORA study. *Depress Anxiety*. 2022;39(1):56-70. doi: [10.1002/da.23223](https://doi.org/10.1002/da.23223).
110. Arora D, Belsiyal CX, Rawat VS. Prevalence and determinants of posttraumatic stress disorder and depression among survivors of motor vehicle accidents from a hilly Indian state. *Indian J Psychiatry*. 2021;63(3):250-7. doi: [10.4103/psychiatry.IndianJPsychiatry_1059_20](https://doi.org/10.4103/psychiatry.IndianJPsychiatry_1059_20).
111. Yrondi A, Colineaux H, Claudet I, Sales de Gauzy J, Huo S, Taib S, et al. Prevalence and prediction of PTSD and depression in mothers of children surviving a motor vehicle crash. *Eur J Psychotraumatol*. 2022;13(2):2121014. doi: [10.1080/2008066.2022.2121014](https://doi.org/10.1080/2008066.2022.2121014).
112. Neylan TC, Kessler RC, Ressler KJ, Clifford G, Beaudoin FL, An X, et al. Prior sleep problems and adverse post-traumatic neuropsychiatric sequelae of motor vehicle collision in the AURORA study. *Sleep*. 2021;44(3):zsaa200. doi: [10.1093/sleep/zsaa200](https://doi.org/10.1093/sleep/zsaa200).
113. Salehi-Pourmehr H, Sabahi Z, Pashazadeh F, Tahmasbi F, Aletaha R, Kashtkar M, et al. Delayed PTSD prevalence in road traffic crashes: a systematic review and meta-analysis. *Curr Psychiatry Res Rev*. 2025;21(3):296-316. doi: [10.2174/0126660822285345240418120620](https://doi.org/10.2174/0126660822285345240418120620).
114. Dai W, Liu A, Kaminga AC, Deng J, Lai Z, Wen SW. Prevalence of posttraumatic stress disorder among children and adolescents following road traffic accidents: a meta-analysis. *Can J Psychiatry*. 2018;63(12):798-808. doi: [10.1177/0706743718792194](https://doi.org/10.1177/0706743718792194).
115. Marasini G, Caleffi F, Machado LM, Pereira BM. Psychological consequences of motor vehicle accidents: a systematic review. *Transp Res Part F Traffic Psychol Behav*. 2022;89:249-64. doi: [10.1016/j.trf.2022.06.017](https://doi.org/10.1016/j.trf.2022.06.017).
116. Kenardy J, Edmed SL, Shourie S, Warren J, Crothers A, Brown EA, et al. Changing patterns in the prevalence of posttraumatic stress disorder, major depressive episode and generalized anxiety disorder over 24 months following a road traffic crash: results from the UQ SuPPORT study. *J Affect Disord*. 2018;236:172-9. doi: [10.1016/j.jad.2018.04.090](https://doi.org/10.1016/j.jad.2018.04.090).
117. Grant DM, Beck JG, Marques L, Palyo SA, Clapp JD. The structure of distress following trauma: posttraumatic stress disorder, major depressive disorder, and generalized anxiety disorder. *J Abnorm Psychol*. 2008;117(3):662-72. doi: [10.1037/a0012591](https://doi.org/10.1037/a0012591).
118. Pozzato I, Craig A, Gopinath B, Kifley A, Tran Y, Jagnoor J, et al. Outcomes after traffic injury: mental health comorbidity and relationship with pain interference. *BMC Psychiatry*. 2020;20(1):189. doi: [10.1186/s12888-020-02601-4](https://doi.org/10.1186/s12888-020-02601-4).
119. Murray J, Ehlers A, Mayou RA. Dissociation and post-traumatic stress disorder: two prospective studies of road traffic accident survivors. *Br J Psychiatry*. 2002;180:363-8. doi: [10.1192/bjp.180.4.363](https://doi.org/10.1192/bjp.180.4.363).
120. Jaspers JP. Whiplash and post-traumatic stress disorder. *Disabil Rehabil*. 1998;20(11):397-404. doi: [10.3109/09638289809166101](https://doi.org/10.3109/09638289809166101).
121. Scherrer JF, Salas J, Norman SB, Schnurr PP, Chard KM, Tuerk P, et al. Association between clinically meaningful posttraumatic stress disorder improvement and risk of type 2 diabetes. *JAMA Psychiatry*. 2019;76(11):1159-66. doi: [10.1001/jamapsychiatry.2019.2096](https://doi.org/10.1001/jamapsychiatry.2019.2096).
122. Shalev AY, Freedman S, Peri T, Brandes D, Sahar T. Predicting PTSD in trauma survivors: prospective evaluation of self-report and clinician-administered instruments. *Br J Psychiatry*. 1997;170:558-64. doi: [10.1192/bjp.170.6.558](https://doi.org/10.1192/bjp.170.6.558).
123. Engelhard IM, van den Hout MA, Schouten EG. Neuroticism and low educational level predict the risk of posttraumatic stress disorder in women after miscarriage or stillbirth. *Gen Hosp Psychiatry*. 2006;28(5):414-7. doi: [10.1016/j.genhosppsych.2006.07.001](https://doi.org/10.1016/j.genhosppsych.2006.07.001).
124. Lunney CA, Schnurr PP, Cook JM. Comparison of clinician- and self-assessments of posttraumatic stress symptoms in older versus younger veterans. *J Trauma Stress*. 2014;27(2):144-51. doi: [10.1002/jts.21908](https://doi.org/10.1002/jts.21908).